

---

# Multivariable Calculus

Math 212 §2 Fall 2014  
©2014 Ron Buckmire

Fowler 309 MWF 11:45am - 12:40pm  
<http://faculty.oxy.edu/ron/math/212/14/>

---

## Class 4: Friday September 5

---

**TITLE** Functions, Vector Functions, Scalar Functions and  $f(x, y)$  as surfaces in  $\mathbb{R}^3$

**CURRENT READING** McCallum, Section 12.1 to 12.2

**HW #2 (DUE WED 09/10/14)**

McCallum, Section 13.3: 2, 5, 6, 10, 20, 22, 29, 35, 38, 81\*.

Section 13.4: 3, 4, 13, 15, 18, 20, 32, 51, 64\*.

Section 17.1: 7, 10, 13, 16, 36, 50\*.

**SUMMARY** In today's class we will begin to learn about functions of two variables, using our intuition from single-variable Calculus to interpret graphs of functions of two variables as surfaces.

---

**DEFINITION: function**

A **function** consists of a pre-image or **domain** (the set of input values), a range or **image** (the set of output values) and a **rule** assigning a unique output value to each input value.

**Exercise**

Write down an example of a function. Explicitly state what the domain, image and rule are for your choice.

### Vector Functions of a Scalar Variable

A **vector** function  $f$  of a **scalar** variable  $\vec{f}(x)$  with **domain**  $D \subset \mathbb{R}$  and **image**  $R \subset \mathbb{R}^n$  means that the function  $f$  has possible input values which form a subset of the real numbers and the set of possible output values are a subset of  $\mathbb{R}^n$ , i.e. vectors. Often the notation  $f : D \rightarrow R$  or  $f : \mathbb{R} \rightarrow \mathbb{R}^n$  is used.

**EXAMPLE**

What kind of geometric object is the image of the function  $\vec{x}(t) = (1 + 3t, -1 - t, -2 + t)$ ?

**NOTE** if the functions in the components of the vector function  $\vec{x}(t)$  are not linear functions of the variable  $t$  (often called the **parameter**), then this 1-dimensional geometric object is called a **parametric curve** in  $\mathbb{R}^n$

## Scalar Functions of a Vector Variable

A **scalar** function  $f$  of a **vector** variable  $f(\vec{x})$  with **domain**  $D \subset \mathbb{R}^n$  and **image**  $R \subset \mathbb{R}$  means that its possible input values are vectors in  $\mathbb{R}^n$  and the set of possible output values are real numbers. Often the notation  $f : D \rightarrow R$  or  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is used.

### DEFINITION: graph

The **graph** of a scalar function of a vector variable  $f(\vec{x})$  is defined to be the set of ordered pairs  $(\vec{x}, f(\vec{x}))$  where  $\vec{x}$  is in the domain of  $f$ . In this case we say that the graph of  $f$  is **explicitly** represented by  $f$ . A graph is a **visual representation** of a function.

**QUESTION:** What are some other ways to represent a function in addition to a graph?

**QUESTION:** Can a function be treated as an object? If so, give an example of this practice!

In practice the only scalar functions of a vector variable that we can really get a good handle on visually are either of the type  $f : \mathbb{R} \rightarrow \mathbb{R}$  or  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ . These graphs are represented by ordered pairs that look like  $(x, f(x))$  and  $(x, y, f(x, y))$  respectively.

### DEFINITION: surface

We know all about the first case from single-variable Calculus so we will be concentrating on the second case, which are often called **surfaces** and denoted  $z = f(x, y)$  so that the ordered pair looks like  $(x, y, z)$ . Below are two examples of surfaces in  $\mathbb{R}^3$ .

