
Multivariable Calculus

Math 212 Spring 2015
©2015 Ron Buckmire

Fowler 309 MWF 9:35am - 10:30am
<http://faculty.oxy.edu/ron/math/212/15/>

Worksheet 13

TITLE Second Order Partial Derivatives

CURRENT READING McCallum, Section 14.7

HW #6 (DUE WEDNESDAY 03/18/15)

McCallum, *Section 14.6*: 4, 11, 12, 26, 34, 35, 47*.

McCallum, *Section 14.7*: 6, 7, 8, 12, 19, 24, 30, 31, 41*.

McCallum, *Section 14.8*: 3, 12, 19*.

McCallum, *Chapter 14 Review*: 2, 14, 35, 45, 64*.

SUMMARY

This worksheet discusses higher order partial derivatives of multivariable functions and introduces the concept of the mixed partial derivative.

RECALL: second derivative

Given an infinitely differentiable function $y = f(x)$ its derivative $f'(x)$ represents the slope of the graph of the function at any point and $f''(x)$ represents the concavity of the graph. Also, $f'(x)$ represents the instantaneous rate of change of $f(x)$ at a point while $f''(x)$ represents the instantaneous rate of change of $f'(x)$.

The Second-Order Partial Derivatives of $f(x, y)$

DEFINITION: f_{xx} , f_{xy} , f_{yy} and f_{yx}

Given a function $z = f(x, y)$ with continuous partial derivatives we can not only find the rate of change with f with respect to x and the rate of change of f with respect to y but the rate of change of *those functions* with respect to x and y also!

$$\frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} \right) = (f_x)_x = f_{xx} \quad \text{and} \quad \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right) = (f_x)_y = f_{xy}$$

$$\frac{\partial}{\partial y} \left(\frac{\partial f}{\partial y} \right) = (f_y)_y = f_{yy} \quad \text{and} \quad \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y} \right) = (f_y)_x = f_{yx}$$

These expressions above are referred to as the second order partial derivatives of $f(x, y)$

EXAMPLE

McCallum, page 812, Exercise 4.

Compute the four second-order partial derivatives of $f(x, y) = e^{2xy}$

QUESTION: Do you notice a relationship between f_{xy} and f_{yx} ?

When Mixed Partial Derivatives Are Equal**THEOREM**

(Clairault's Theorem) If f_{yx} and f_{xy} are continuous at some point (a, b) found in a disc $(x - a)^2 + (y - b)^2 \leq D$ for some $D > 0$ on which $f(x, y)$ is defined, then $f_{xy}(a, b) = f_{yx}(a, b)$.

Applications of the Second-Order Partial Derivatives

Recall (from *Worksheet #8*) that the local linearization of a function $f(x, y)$ near the point (a, b) is given by the tangent plane

$$f(x, y) \approx P(x, y) = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b) \quad (1)$$

Taylor Polynomial Approximations

Note that the expression on the right hand side of (1) can be thought of as Taylor Polynomial of Degree 1 approximating $f(x, y)$ near (a, b) for a function that has continuous first-order partial derivatives.

We can expand this idea from (1) to improve our approximation of this function. If $f(x, y)$ has continuous second-order partial derivatives we can produce a Taylor Polynomial of Degree 2 approximating $f(x, y)$ near (a, b) :

$$f(x, y) \approx Q(x, y) = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b) + \frac{f_{xx}(a, b)}{2}(x - a)^2 + f_{xy}(a, b)(x - a)(y - b) + \frac{f_{yy}(a, b)}{2}(y - b)^2$$

Exercise

McCallum, page 811, Example 5.

Find the Taylor Polynomial of degree 2 at the point $(1, 2)$ for the function $f(x, y) = \frac{1}{xy}$.


GROUPWORK

You are told that there is a function f whose partial derivative $f_x(x, y) = x + 4y$ and $f_y(x, y) = 3x - y$. Do you believe this? PROVE YOUR ANSWER!

The kinetic energy of a body with mass m and velocity v is $K = \frac{1}{2}mv^2$. Show that $\frac{\partial K}{\partial m} \frac{\partial^2 K}{\partial v^2} = K$.

The gas law for fixed mass m of an ideal gas at the absolute temperature T , pressure P and volume V is $PV = mRT$ where R is the gas constant. Show that

$$\frac{\partial P}{\partial V} \frac{\partial V}{\partial T} \frac{\partial T}{\partial P} = -1$$

**Exercise**

Consider the figure with the contour diagram of an unknown function $f(x, y)$. Estimate f_{xx} , f_{yy} and f_{xy} at the indicated points.