

---

# Linear Systems

Math 214 Spring 2006  
©2006 Ron Buckmire

Fowler 307 MWF 2:30pm - 3:25pm  
<http://faculty.oxy.edu/ron/math/214/06/>

---

*Class 2: Wednesday January 25*

**SUMMARY** Lengths and Dot Products

**CURRENT READING** Poole 1.2

## RECALL

Previously we have discussed addition and scalar multiplication of vectors, primarily in the form of linear combinations of vectors. Today we're going to think about *multiplication* of vectors. As with most topics in this course, there's an algebraic view and a geometric (graphical) view of understanding this concept.

---

---

*Homework Assignment #2*

*Section 1.2 # 2, 5, 11, 17, 19, 25, 44, 46, 47, 52: DUE FRI JAN 27*

---

---

Consider the vectors  $\vec{v} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$  and  $\vec{w} = \begin{bmatrix} w_1 \\ w_2 \end{bmatrix}$

### 1. The Dot Product

$$\vec{v} \cdot \vec{w} = v_1 w_1 + v_2 w_2$$

Note this product is a scalar, not a vector. The dot product operation IS commutative. Can you PROVE this?

Also, interestingly  $\vec{v} \cdot \vec{v} = v_1 v_1 + v_2 v_2 = v_1^2 + v_2^2$

the above formula should remind you of the expression for the length or magnitude of a vector  $\vec{v}$ , which is usually denoted  $\|\vec{v}\|$

So,

$$\|\vec{v}\| = \sqrt{\vec{v} \cdot \vec{v}}$$

### 2. Normalization

Oftentimes we want to work with vectors of unit length. These vectors are called *normalized*.

Suppose  $\vec{w} = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$  How can we normalize  $\vec{w}$  so that it has the same direction, but magnitude 1?

In general, a vector  $\vec{v}$  is normalized by \_\_\_\_\_.

### 3. Orthogonality.

Suppose  $\vec{a} \cdot \vec{b} = 0$  what can we say about  $\vec{a}$  and  $\vec{b}$ ?

$$\text{Consider } \vec{a} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \vec{b} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

AND

$$\vec{a} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}, \vec{b} = \begin{bmatrix} -6 \\ 4 \end{bmatrix}$$

AND

$$\vec{a} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \vec{b} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

Do you notice a pattern among these vectors? What properties do the vector pairs share?

### 4. Angle Between Vectors.

$$\text{Consider the vectors } \vec{v} = \begin{bmatrix} \cos \alpha \\ \sin \alpha \end{bmatrix} \text{ and } \vec{w} = \begin{bmatrix} \cos \beta \\ \sin \beta \end{bmatrix}$$

$$\vec{v} \cdot \vec{w} = \underline{\hspace{10em}}$$

Can you rewrite this formula using a trigonometric identity?

$$\text{Draw } \vec{a} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \vec{b} = \begin{bmatrix} -\sqrt{2}/2 \\ \sqrt{2}/2 \end{bmatrix} \text{ on an axes.}$$

1. What is the dot product between them?
2. What is the angle between them?
3. What is the angle between  $\vec{a}$  and  $2\vec{b}$ ?

5. Angle Formula. For any vectors  $\vec{v}$  and  $\vec{w}$  where  $\theta$  is the smaller angle between them,

$$\cos \theta = \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|}$$

How can we use this formula to confirm our understanding of orthogonality? (i.e. what happens when the dot product between two vectors is zero?)

6. **Projection.** For any vectors  $\vec{u}$  and  $\vec{v}$  where  $\vec{u} \neq 0$  then **the projection of  $\vec{v}$  onto  $\vec{u}$**  is the vector  $\text{proj}_{\vec{u}}(\vec{v})$  defined by:

$$\text{proj}_{\vec{u}}(\vec{v}) = \left( \frac{\vec{u} \cdot \vec{v}}{\vec{u} \cdot \vec{u}} \right) \vec{u}$$