

---

# Linear Systems

Math 214 Spring 2007  
©2007 Ron Buckmire

Fowler 110 MWF 2:30pm - 3:25pm  
<http://faculty.oxy.edu/ron/math/214/07/>

---

*Class 7: Wednesday February 7*

**SUMMARY** Reduced Row Echelon Form and Rank

**CURRENT READING** Poole 2.2

## Summary

We will discuss the reduced row echelon form of a matrix, sometimes denoted  $\mathbf{rref}(A)$ , and introduce the important concept of rank of a matrix.

---

---

## Homework Assignment

*HW #7: Section 2.2: 1, 2, 3, 4, 5, 6, 7, 8, 22, 23, 26, 27, 28, 29, 36, 43. EXTRA CREDIT # 47.*

---

---

### 1. Reduced Row Echelon Form

#### Definition

A matrix is in **reduced row echelon form** if it satisfies the following property:

1. It is in row echelon form (i.e. all zero rows are at the bottom and the first non-zero entry of each non-zero row is in a column to the left of any non-zero leading entry in rows below it, forming an echelon or staircase appearance).
2. The leading entry is in each non-zero row is 1.
3. Each column containing a leading entry of 1 has zeros everywhere else.

#### GROUPWORK

Find the reduced row echelon matrix for each of the following. (Also note what are the dimensions (i.e. number of rows and number of columns) for each matrix?)

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 3 \\ 2 & 6 \end{bmatrix}$$

$$B = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & -1 \end{bmatrix}$$

$$C = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & -1 \\ 0 & 4 & 8 \end{bmatrix}$$

## 2. Definition of Rank

*Definition 1.* The rank of a  $m \times n$  matrix  $A$  is the number of non-zero rows in its row echelon form. We call this number  $r = \text{rank}(A)$ .

By definition  $r \leq m$  and  $r \leq n$ . Also,  $r$  is the number of pivots a coefficient matrix has when applying Gaussian Elimination.

### **THEOREM**

When there are more columns than rows in a matrix, i.e.  $n > m$  there is ALWAYS atleast one free variable. In other words if  $A$  is the  $m \times n$  coefficient matrix of a consistent linear system, the number of free variables =  $n - r$ . This result is known as **the Rank Theorem**.

### **Exercise**

What are the rank of each of the matrices  $A$ ,  $B$  and  $C$  given on the other side of this page? How is  $r$  related to  $m$  and  $n$  in each case? Do you notice a pattern?

### **EXAMPLE**

Let's solve the following linear system of equations

$$\begin{aligned}x + 2y - z &= 3 \\2x + 3y + z &= 1\end{aligned}$$

### **EXAMPLE**

Consider  $\vec{p} = (1, 0, -1)$ ,  $\vec{q} = (0, 2, 1)$ ,  $\vec{u} = (1, 1, 1)$  and  $\vec{v} = (3, -1, -1)$  Do the following lines  $\vec{x} = \vec{p} + t\vec{q}$  and  $\vec{x} = \vec{u} + t\vec{v}$  intersect? If so, where? Is it possible for lines in  $\mathbb{R}^3$  to not intersect and not be parallel?