## Complex Analysis

Math 312 Fall 2001 ©Buckmire M 5-6:25, R 1:30-2:55 Fowler 112, Fowler 201

TEST 1: Friday October 19, 2001

**Directions**: Read *all* 4 problems first before answering any. **You may choose to answer question 3 or question 4.** You may have access to any notes or the textbook. This is a one hour test. You have 90 minutes to complete it.

No.	Score	Maximum
1		40
2		30
3 or 4		30
Total		100

- 1. [40 pts. total] Mapping. We want to find the general form of the function M(z) = Az + B which maps one circle, the set of points  $\mathbf{P}$ :  $|z z_0| = r$ , to another circle located somewhere else, the set of points  $\mathbf{Q}$ :  $|w w_0| = \rho$  in the complex plane.
- (a) [5 pts] Find a mapping of the form  $f_1(z) = \alpha z$  which maps **P** so that it has the same radius as **Q**.

(b) [5 pts] Find a mapping of the form  $f_2(z) = z + \beta$  which maps **P** so that its center is at the same location as **Q**.

(c) [10 pts] Will  $f_2(f_1(z)) = F(z)$  be the mapping which maps **P** to **Q**? In other words, what is the image of **P** under F(z)?

(d) [10 pts] Find an example of a mapping M(z) = Az + B where A and B depend on the parameters r,  $\rho$ ,  $z_0$  and  $w_0$  which maps **P** to **Q**.

(e) [10 pts] Use your answers above to find the function M(z) which maps |z-2-i|=1 to |w+2+3i|=2.

**2.** [30 pts.] **Arithmetic of Complex Numbers.** (a) [10 pts] What condition on a and b must be met for  $(a + bi)^2 = ci$  where a, b and c are all real numbers? Where in the complex plane would (a, b) have to be for c to be negative?

**(b)** [10 pts] Use your answer in part **(a)** to help you evaluate  $\sqrt{8i}$ .

(c) [10 pts] Use your answers above to find all the solutions of  $z^2 + 2iz + 8i - 1 = 0$ .

- 3. [30 pts. total] Cauchy-Riemann Equations, Harmonic Functions. Consider the function  $f = u(x, y) + iv(x, y) = x^2 + y^2 + 2xyi$
- (a) [10 pts] Show that the set of points for which f'(z) exists all lie on the x-axis.

(b) [10 pts] Using your information from (a), on what set of points is f(z) analytic? Explain your answer.

(c) [10 pts] Show that v(x,y) is harmonic. Is the given u(x,y) its harmonic conjugate? If not, find the harmonic conjugate of v(x,y)

4. [30 pts. total] Analyticity, Differentiability.

The **Jacobian** of a mapping u = u(x, y), v = v(x, y) from the xy-plane to the uv-plane is defined to be the determinant

$$J(x_0,y_0) = \left| egin{array}{cc} rac{\partial u}{\partial x} & rac{\partial u}{\partial y} \ rac{\partial v}{\partial x} & rac{\partial v}{\partial y} \end{array} 
ight|,$$

where the partial derivatives  $u_x$ ,  $u_y$ ,  $v_x$ ,  $v_y$  are all evaluated at  $(x_0, y_0)$ .

(a) [10 pts] If f = u + iv is analytic on a neighborhood containing  $z_0 = x_0 + iy_0$  show that  $J(x_0, y_0) = |f'(z_0)|^2$ .

**(b)** [20 pts] For the function f(z) = Az + B find J(0,0) two different ways (i.e. from the definition and from the result given in part (a).