
Complex Analysis

Math 312 Spring 1998
© Buckmire

MWF 10:30am - 11:25am
Fowler 112

Class 28 (Friday April 3)

SUMMARY Poles, Zeroes, Residues and Cauchy's Residue Theorem

CURRENT READING Brown & Churchill pages 193-196

NEXT READING Brown & Churchill pages 180-196

Zeroes and Poles

So, we have had a lot of experience finding the **poles** of a function. The problem of finding a pole is equivalent to finding the **zero** of a related function. Let's formalize these definitions:

Zero

A point z_0 is called a **zero of order m** for the function $f(z)$ if f is analytic at z_0 and f and its first $m - 1$ derivatives vanish at z_0 , but $f^{(m)}(z_0) \neq 0$.

Pole

A point z_0 is called a **pole of order m** of $f(z)$ if $1/f$ has a zero of order m at z_0 .

Identifying Poles and Zeroes

Let f be analytic. Then f has a **zero of order m** at z_0 if and only if $f(z)$ can be written as $f(z) = g(z)(z - z_0)^m$ where g is analytic at z_0 and $g(z_0) \neq 0$.

If $f(z)$ can be written as $f(z) = \frac{g(z)}{(z - z_0)^m}$ where $g(z)$ is analytic at z_0 , then f has a **pole of order m** at $z = z_0$ and $g(z_0) \neq 0$

How do we find poles of a function? Well, if we have a quotient function $f(z) = p(z)/q(z)$ where $p(z)$ and $q(z)$ are analytic at z_0 and $p(z_0) \neq 0$ then $f(z)$ has a pole of order m if and only if $q(z)$ has a zero of order m .

Exercise

We will classify all the singularities of $f(z) = \frac{3z + 2}{z^4 + 1}$

Groupwork

Let's try and classify all the singularities of the following functions:

(a) $f(z) = \frac{\cos z}{z^2(z - \pi)^3}$

(b) $f(z) = \frac{\sin z}{z^2 - 4}$

(c) $f(z) = \tan z$

(d) $f(z) = \frac{z}{z^2 - 6z + 10}$

Residues

Once we know all the singularities of a function it is useful to compute the residues of that function. If a function $f(z)$ has a pole of order m at z_0 , the residue, denoted by $\mathbf{Res}(f; z_0)$ or $\mathbf{Res}(z_0)$ is given by the formula below:

$$\mathbf{Res}(f; z_0) = \lim_{z \rightarrow z_0} \left\{ \frac{1}{(m-1)!} \frac{d^{m-1}}{dz^{m-1}} [(z-z_0)^m f(z)] \right\}$$

Find the **residues** of all the singularities we previously classified.

Cauchy's Residue Theorem

If f is analytic on a simple (positively oriented) closed contour and everywhere inside except the finite number of points z_1, z_2, \dots, z_n inside, then

$$\oint f(z) dz = 2\pi i \sum_{k=1}^n \mathbf{Res}(z_k)$$

Use Cauchy Residue Theorem to evaluate the following integrals:

(a) $\oint_{|z|=5} \frac{\cos z}{z^2(z-\pi)^3} dz$

(b) $\oint_{|z|=5\pi} \frac{\sin z}{z^2-4} dz$

(c) $\oint_{|z|=2\pi} \tan z dz$

(d) $\oint_{|z|=8} \frac{z}{z^2-6z+10} dz$

(e) $\oint_{|z|=2} \frac{3z+2}{z^2+1} dz$