

---

# Differential Equations

Math 341 Fall 2010

MWF 2:30-3:25pm Fowler 307

©2010 Ron Buckmire

<http://faculty.oxy.edu/ron/math/341/10/>

---

## Worksheet 8: Monday September 28

**TITLE** Linear Differential Equations

**CURRENT READING** Blanchard, 1.8

---

### Homework Assignments due Friday September 24

Section 1.7: 3, 6, 8, 12, 15.

Section 1.8: 4, 5, 8, 9, 17, 18, 20. Section 1.9: 4, 5, 9, 12, 19.

---

### SUMMARY

We will learn about the Linearity Principle and how it helps us to solve an entire class of ODEs. In addition we will learn some important new terms to describe ODEs, **homogeneous** and **non-homogeneous**. We'll try and make some analogies to some ideas already seen in Linear Algebra.

---

### 1. Linearity Principles

#### DEFINITION: first-order linear DE

A **first-order linear DE** has the form  $\frac{dy}{dt} = a(t)y + b(t)$ . When  $b(t) = 0$  for all  $t$  the equation is called **homogeneous**, otherwise the DE is called **non-homogeneous**.

The operator form of a nonhomogeneous first-order linear ODE is

$$\begin{aligned}\frac{dy}{dt} - a(t)y &= b(t) \\ \left[ \frac{d}{dt} - a(t) \right] y &= b(t) \\ \mathcal{L}y &= b(t)\end{aligned}$$

The operator  $\mathcal{L}$  represents what actions are applied to the function  $y(t)$ . An operator is a mathematical object which takes a function as an input and output. Clearly, if  $b(t) = 0$  then  $\mathcal{L}y = 0$  and we name solutions such equations as homogeneous solutions and denote them  $y_h$ . A solution which solves  $\mathcal{L}y = b$  is denoted  $y_p$  and called a particular solution.

Differential Operators obey linearity properties reminiscent of Linear Transformations from Linear Algebra (Math 214). In other words, given any two function  $y_1$  and  $y_2$  and a constant  $c \in \mathbb{R}$ ,

$$\mathcal{L}(y_1 + y_2) = \mathcal{L}y_1 + \mathcal{L}y_2 \text{ and } \mathcal{L}(cy) = c\mathcal{L}y$$

### RECALL

A transformation (or mapping or function)  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is called a **linear transformation** if

1.  $T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$  for all  $\vec{u}$  and  $\vec{v}$  in  $\mathbb{R}^n$
2.  $T(c\vec{v}) = cT(\vec{v})$  for all  $\vec{v}$  in  $\mathbb{R}^n$  and all scalars  $c$ .

**THEOREM**

If  $y_h(t)$  is a solution of the homogeneous DE then so is  $cy(t)$  where  $c$  is any real number.  
If  $y_1(t)$  and  $y_2(t)$  are both solutions of the homogeneous DE, then so is  $y_1(t) + y_2(t)$ .

**Exercise**

Prove either one of the above statements, using operator notation.

**THEOREM: Extended Linearity Principles**

- 1) If  $y_h(t)$  is any solution of the homogeneous DE and  $y_p(t)$  is any solution of the non-homogeneous DE then  $y_h + y_p$  is a solution of the non-homogeneous DE.
- 2) If  $y_1(t)$  and  $y_2(t)$  are solutions of the non-homogeneous DE, then  $y_1(t) - y_2(t)$  is a solution of the associated homogeneous DE.

**PROOF**

Together we'll prove both of the above statements, using operator notation.

**2. General Solution to a Linear ODE**

The implication of the extended linearity principle is the idea that the general solution  $y_g(t)$  to a non-homogeneous DE can be written as the sum of the general solution of the homogeneous DE  $y_h(t)$  and one solution of the non-homogeneous DE  $y_p(t)$ . This is an incredibly important idea that repeats itself throughout the field of differential equations and other branches of mathematics.

This idea is very similar to the idea in Linear Algebra that the general solution to the linear system problem  $A\vec{x} = \vec{b}$  is a vector  $\vec{x}$  that can be written as a sum of two vectors,  $\vec{x} = \vec{x}_h + \vec{x}_p$ , where  $\vec{x}_h$  is in the nullspace of  $A$  (i.e. solves the homogeneous problem  $A\vec{x}_h = \vec{0}$ ) and  $\vec{x}_p$  is in the rowspace of  $A$  (i.e. a vector which satisfies  $A\vec{x}_p = \vec{b}$ ).

**EXAMPLE**

Let's try to find the general solution of the linear DE  $\frac{dy}{dt} = -2y + e^t$ .

**GROUPWORK**

Can we use the same technique to solve  $\frac{dy}{dt} = -2y + 3e^{-2t}$ ? Let's try.

**3. The Method of Undetermined Coefficients OR How To Guess Smart** Given a linear DE of the form  $\mathcal{L}y = f$  depending on what kind of function  $f$  is we can make various intelligent guesses as to what the particular form of the solution  $y_p$  will take.

We need to determine what “family of functions” the right hand side function belongs to, as this determines what our intelligent guess will be. The Table below should give you an example of most functions that you are likely to run into.

Function	Family	Guess
$k$ (a constant)	1	A
$x^n$ ( $n$ integer)	Polynomial	$A + Bx + Cx^2 + \dots Qx^n$
$e^{kx}$ ( $k$ constant)	Exponential	$Ae^{kx}$
$\sin(kx)$ ( $k$ constant)	Trigonometric	$A \sin(kx) + B \cos(kx)$
$\cos(kx)$ ( $k$ constant)	Trigonometric	$A \sin(kx) + B \cos(kx)$

**EXAMPLE**

Blanchard, page 125, #20.

What would a good guess for the solution to  $\frac{dy}{dt} + 2y = 3t^2 + 2t - 1$  be?

**Homework**

Blanchard, page 123, #4.

Find the general solution of  $\frac{dy}{dt} = 2y + \sin(2t)$ .