\mathbf{D} ifferential \mathbf{E} quations

Math 341 Fall 2013 © 2013 Ron Buckmire

 $MWF~12:50-1:45pm~Fowler~307 \\ \texttt{http://faculty.oxy.edu/ron/math/341/13/}$

Class 5: Monday September 9

TITLE Existence and Uniqueness of Solutions

CURRENT READING Blanchard, 1.5

Homework Set #3 due Friday September 20

Section 1.4: 2, 6, 11, 15.

Section 1.5: 2, 3, 12, 14, 15.

Section 1.6: 2, 7, 8, 19, 20, 30, 31, 41.

SUMMARY

We will investigate the conditions which guarantee existence and/or uniqueness of solutions to the initial value problem y' = f(t, y), $y(t_0) = y_0$.

1. Do Problems Always Have Solutions?

Think about the equation $2x^5 - 10x + 5 = 0$. Does it have a solution? How do we know?

2. Existence and Uniqueness Of Particular Solutions

The main questions we would like to be able to answer when analyzing IVPs are:

- 1) **Existence** Does the differential equation possess solutions which pass through the given initial condition? and
- 2) Uniqueness If such a solution does exist, can we be certain that it is the only one? Luckily, there's a theorem that answers these questions for us.

THEOREM: Existence of a Unique Solution

Let \mathcal{R} be a rectangular region in the xy-plane defined by $a \leq x \leq b, c \leq y \leq d$ that contains the point (x_0, y_0) in its interior. IF f(x, y) and $\partial f/\partial y$ are continuous on \mathcal{R} , THEN there exists some interval I_0 defined as $x_0 - h < x < x_0 + h$ for h > 0 contained in $a \leq x \leq b$ and a unique function y(x) defined on I_0 that is a solution of the initial value problem y' = f(x, y) $y(x_0) = y_0$.

3. Theorems Have Hypotheses and Conclusions

The existence and uniqueness theorem is actually two different theorems with different hypotheses and conclusions.

- 1) **Existence** IF f(t, y) is continuous on a square containing t_0, y_0 THEN there exists a solution on an interval $(t_0 \epsilon, t_0 + \epsilon)$ for some $\epsilon > 0$
- 2) Uniqueness IF f(t,y) and $\frac{\partial f}{\partial y}$ are both continuous on a square containing (t_0, y_0) THEN there exists a unique solution on an interval $(t_0 \epsilon, t_0 + \epsilon)$ for some $\epsilon > 0$

RECALL

IF A, THEN B is equivalent to (The Contrapositive) IF NOT B, THEN NOT A.

(The Inverse) IF NOT A, THEN NOT B and (The Converse) IF B, THEN A are equivalent to each other, but are NOT equivalent to the original theorem $A \Rightarrow B$.

For example consid	er the logical statement:	"IF it is raining,	THEN the g	ground is wet."
The contrapositive	is "If the ground is NOT	wet, then it is N	OT raining."	,
The Converse: IF		THEN		
(T) T TD		TITINI		

EXAMPLE

Show that the initial value problem

$$\frac{dy}{dx} = x\sqrt{y}, \quad y(0) = 0$$

has at least two solutions since the trivial solution y(x) = 0 and the solution $y(x) = \frac{1}{16}x^4$ both satisfy the IVP. **Verify this!**

Using the Existence and Uniqueness Theorem, we look at the functions $f(x,y) = x\sqrt{y}$ and $\frac{\partial f}{\partial y} = \frac{x}{2\sqrt{y}}$. At the origin (0,0) what can we say about f(x,y) and $f_y(x,y)$?

What can we say about f(x, y) and $f_y(x, y)$ at (2, 4)? What does this imply about existence and uniqueness of the corresponding IVP $y' = xy^{1/2}, y(2) = 4$?

4. Implications of Existence/Uniqueness Theorem

Exercise

Inspired by Blanchard, Devaney & Hall, #9, page 72.

- (a) Show that $y_1(t) = t^2$ and $y_2(t) = t^2 + 1$ are both solutions of $\frac{dy}{dt} = -y^2 + y + 2yt^2 + 2t t^2 t^4$
- (b) Show that if y(t) is another solution to the given ODE with initial condition 0 < y(0) < 1 then $t^2 < y(t) < t^2 + 1$ for all t
- (c) Illustrate your answer by using technology to explore the slope field

EXAMPLE

Inspired by Blanchard, Devaney & Hall, #16, page 73.

Consider the differential equation $\frac{dy}{dt} = \frac{t}{y-2}$, y(-1) = 0

- (a) Find a formula for the solution to the IVP.
- (b) State the domain of definition of the solution
- (c) Describe what happens as the solution reaches the limit of its domain of definition