
Differential Equations

Math 341 Fall 2014
©2014 Ron Buckmire

MWF 3:00-3:55pm Fowler 307
<http://faculty.oxy.edu/ron/math/341/14/>

Worksheet 26

TITLE The Laplace Transform and The Heaviside Function

CURRENT READING Blanchard, 6.2

Homework #11 Assignments due Monday November 17

Section 6.1: 2, 3, 5, 7, 8, 9, 15, 18, 25*.

Section 6.2: 1, 2, 4, 8, 15, 16, 18*.

Homework #12 Assignments due Monday November 24

Section 6.3: 5, 6, 8, 15, 18, 27, 28.

Section 6.4: 1, 2, 6, 7*.

SUMMARY

We shall continue our analysis of Laplace Transforms by considering discontinuous functions.

1. Translation in t

DEFINITION: Heaviside function

The **unit step function** or **Heaviside function** $\mathcal{H}(t)$ is defined to be **0** when its argument is less than zero and **1** when its argument is greater than or equal to zero. Generally, it is written as \mathcal{H}_a or $\mathcal{H}(t - a) = \begin{cases} 0, & 0 \leq t < a \\ 1, & t \geq a \end{cases}$

NOTE: **Blanchard, Devaney & Hall** uses the notation $u_a(t)$ for $\mathcal{H}(t - a)$.

Exercise

Sketch a picture of $u_a(t)$ in the space below. Is this function piecewise continuous? (Why do we care?) What is the definition of “piecewise continuous”?

Let's show that $\mathcal{L}[\mathcal{H}(t - a)] = \frac{e^{-as}}{s}$

GROUPWORK

Confirm that $f_1(t) = \begin{cases} g(t), & 0 \leq t < a \\ h(t), & t \geq a \end{cases}$

can be written as $f_1(t) = g(t) - g(t)\mathcal{H}(t-a) + h(t)\mathcal{H}(t-a)$ or $f_1(t) = g(t) + \mathcal{H}_a(t)(h(t) - g(t))$

How would you combine Heaviside functions to represent the following function? [HINT: what would the graph of the difference of two Heaviside functions look like?]

$$f_2(t) = \begin{cases} 0, & 0 \leq t < a \\ g(t), & a \leq t < b \\ 0, & t \geq b \end{cases}$$

This kind of function $f_2(t)$ is an example of an **interval function**, and is denoted $u_{ab}(t)$. $u_{ab}(t) = 1$ if $a < t < b$ and 0 otherwise.

EXAMPLE

Blanchard, Devaney & Hall, page 586, #15. Suppose $a \geq 0$. Find the general solution of $\frac{dy}{dt} = -y + u_a(t)$

THEOREM: Second Translation Theorem

If $F(s) = \mathcal{L}[f(t)]$ and $a > 0$ is any positive real number, then $\mathcal{L}[f(t-a)\mathcal{H}(t-a)] = e^{-as}F(s)$.

It directly follows then that $\mathcal{L}[\mathcal{H}(t-a)] = \frac{e^{-as}}{s}$.

Corollary

$$\mathcal{L}^{-1}[e^{-as}F(s)] = f(t-a)\mathcal{H}(t-a)$$

THEOREM: Alternate form of the Second Translation Theorem

It can be annoying to try and get the function which is multiplying the Heaviside function into the form $f(t-a)$ for use in the previous version of the Second Translation Theorem so a more useful results is: $\mathcal{L}[g(t)\mathcal{H}(t-a)] = e^{-as}\mathcal{L}[g(t+a)]$

2. Translation in s

THEOREM: First Translation Theorem

If $F(s) = \mathcal{L}[f(t)]$ and a is any real number, then $\mathcal{L}[e^{at}f(t)] = F(s - a)$. Sometimes the notation $\mathcal{L}[e^{at}f(t)] = \mathcal{L}[f(t)]|_{s \rightarrow s-a}$ is used.

Corollary

The inverse of the First Translation Theorem can be written as $\mathcal{L}^{-1}[F(s - a)] = e^{at}f(t)$.

Exercise Given that $\frac{2s + 5}{(s - 3)^2} = \frac{2}{s - 3} + \frac{11}{(s - 3)^2}$, compute $\mathcal{L}^{-1}\left[\frac{2s + 5}{(s - 3)^2}\right]$. (HINT: recall that $\mathcal{L}^{-1}\left[\frac{1}{s^2}\right] = t$)

EXAMPLE Compute $\mathcal{L}^{-1}\left[\frac{s/2 + 5/3}{s^2 + 4s + 6}\right]$.

HINT: recall $\mathcal{L}^{-1}\left[\frac{s}{s^2 + k^2}\right] = \cos(kt)$ and $\mathcal{L}^{-1}\left[\frac{k}{s^2 + k^2}\right] = \sin(kt)$

EXAMPLE **Zill, Example 3, page 295.** Let's use Laplace Transforms to show that the solution of $y'' - 6y' + 9y = t^2e^{3t}$, $y(0) = 2$, $y'(0) = 17$ is $y(t) = 2e^{3t} + 11te^{3t} + \frac{1}{12}t^4e^{3t}$.