

FISHERIES INVENTORY AND UTILIZATION
OF SAN DIEGO BAY, SAN DIEGO, CALIFORNIA
FOR SURVEYS CONDUCTED IN APRIL AND JULY 2019



Jonathan P. Williams, M.S.

Chelsea M. Williams, A.B.

Zoe Scholz, B.S., B.A.

Matthew J. Robart, M.S.

Daniel J. Pondella, II, M.A., Ph.D

Vantuna Research Group
Moore Laboratory of Zoology
Occidental College
1600 Campus Rd.
Los Angeles, CA 90041

September 2019

Vantuna Research Group

Daniel J. Pondella, II, Ph.D., Principal Investigator
Jonathan P. Williams, M.S., Research Scientist
Chelsea M. Williams, A.B., Research Associate
Matthew J. Robart, M.S., Research Associate
Zoe Scholz, B.S., B.A., Research Assistant

Field and laboratory research assistants:

Jeremy Claisse, Ph.D.
Adam Obaza, M.S.
Bill Power, M.S.
Benjamin Grime, B.A.
Stuart Schwab, B.A.
Sam Soule, B.A.
Aaron Sugimoto, B.A.
Jacob Eagleton, B.S.
Austin Pyles, B.S.
Shahar Amitay
Sarah Ashe
Hannah Hoefs
Kevin Fistanic
Corrie McKee
Clay Pollock
Alayna Schwartz
Xavier Nelson-Rountree

Volunteers:

Ralph Appy (Port of Los Angeles – retired)
Bruno Passarelli (UCLA)
Jake Feingold (University of Chicago)
Heather Kramp (San Diego Unified Port District)
Madelyn Roycroft (San Diego Unified Port District)

Report Authors: Jonathan Williams, Chelsea Williams, Zoe Scholz, Matt Robart, and Dan Pondella

Photographs: All photos were taken by Jonathan Williams, Chelsea Williams, Matt Robart, Clay Pollock, Kevin Fistanic, and Zoe Scholz

Acknowledgments:

We greatly appreciate Eileen Maher's oversight and management of this project, the assistance of Jessica Bredvik and Dana Schrimpf in granting of access to Naval Air Station North Island by the United States Navy, as well as the assistance of all of the field work volunteers listed above.

TABLE OF CONTENTS

List of Tables	iv
List of Figures	v
1.0 Executive Summary	1
2.0 Introduction	4
3.0 Methods and Materials	
3.1 Survey Locations	5
3.2 Ichthyofaunal Sampling	6
3.3 Water Quality Sampling	11
3.4 Data Analysis	
3.4.1 Best Estimate of Density and Standing Stock	12
3.4.2 Community Analysis	13
4.0 Results and Discussion	
4.1 Water Quality Parameters	14
4.2 Numerical Catch and Biomass	16
4.3 Species Diversity and Richness	24
4.4 Catch by Sampling Period and Ecoregion	25
4.5 Catch by Depth Strata and Subhabitats	29
4.6 Nursery Area Function	32
4.7 Ecological Importance of Species	33
4.8 Principal Species	35
4.9 Catch by Sampling Method	45
4.10 Best Estimates of Density and Standing Stock	52
4.11 Avian Forage Species	55
4.12 Fisheries Species	57
4.13 Southern Species Found in San Diego Bay	58
4.14 Indigenous Bay and Estuary Fishes	59
4.15 Invasive Species	60
4.16 Comparison of the Current and Historical April and July Surveys	
4.16.1 Abundance, Biomass, and Stock Estimates	61
4.16.2 Community Metrics	65
4.16.3 Community Structure	68
5.0 Literature Cited	70

LIST OF TABLES

Table 1.	Lambert coordinates for San Diego Bay fisheries inventory	6
Table 2.	Estimates of aerial coverage of depth strata by ecoregion	12
Table 3.	Total abundance of fishes collected in San Diego Bay	18
Table 4.	Total biomass of fishes collected in San Diego Bay	19
Table 5.	Total abundance and biomass in the North Ecoregion	20
Table 6.	Total abundance and biomass in the North-Central Ecoregion	21
Table 7.	Total abundance and biomass in the South-Central Ecoregion	22
Table 8.	Total abundance and biomass in the South Ecoregion	23
Table 9.	Total abundance by sampling period	27
Table 10.	Total biomass by sampling period	28
Table 11.	Total abundance of fishes by depth strata	30
Table 12.	Total abundance by subhabitat	31
Table 13.	Percent of juveniles by species	32
Table 14.	Ecological Index of fishes	34
Table 15.	Total catch by sampling method	46
Table 16.	Total biomass by sampling method	49
Table 17.	Numerical and biomass density estimates by gear type	52
Table 18.	Best estimate of fish density and estimated stock size	53
Table 19.	Best estimate of biomass density and standing stock	54
Table 20.	Best estimate of avian forage fish species	56
Table 21.	Best estimate of fishery species	57
Table 22.	Panamic species from San Diego Bay	58
Table 23.	Abundance of Panamic species in San Diego Bay	59
Table 24.	Indigenous bay/estuarine species taken in San Diego Bay	59
Table 25.	Historical comparison of total catches	61
Table 26.	Historical comparison of abundance for top species	62
Table 27.	Historical comparison of total biomass	63
Table 28.	Historical comparison of biomass for top species	64
Table 29.	Historical comparison of stock/biomass standing stock estimates	65
Table 30.	Shannon diversity historical comparison	66
Table 31.	Species richness historical comparison	67

LIST OF FIGURES

Figure 1.	Sampling locations in San Diego Bay	5
Figure 2.	Sampling events for the North Ecoregion	8
Figure 3.	Sampling events for the North-Central Ecoregion	9
Figure 4.	Sampling events for the South-Central Ecoregion	10
Figure 5.	Sampling events for the South Ecoregion	11
Figure 6.	Summary of physical-chemical measurements by ecoregion	14
Figure 7.	Monthly rainfall in San Diego	15
Figure 8.	Catch of fishes by Ecoregion	16
Figure 9.	Catch of the five numerically dominant species by ecoregion	17
Figure 10.	Shannon diversity and species richness by ecoregion	24
Figure 11.	Shannon diversity and richness by sampling month	24
Figure 12.	Total abundance by sampling month	26
Figure 13.	Total biomass by sampling month	26
Figure 14.	Top 20 species ranked by Ecological Index	33
Figure 15.	Length frequency of Topsmelt	35
Figure 16.	Disc width frequency of Round Stingray	36
Figure 17.	Length frequency of Spotted Sand Bass	37
Figure 18.	Length frequency of Slough Anchovy	38
Figure 19.	Length frequency of Deepbody Anchovy	39
Figure 20.	Length frequency of Giant Kelpfish	40
Figure 21.	Length frequency of Shiner Perch	41
Figure 22.	Length frequency of Kelp Pipefish	42
Figure 23.	Length frequency of California Halibut	43
Figure 24.	Length frequency of Dwarf Perch	44
Figure 25.	Historical comparison of total catch	61
Figure 26.	Historical comparison of abundance for top species	62
Figure 27.	Historical comparison of total biomass	63
Figure 28.	Historical comparison of biomass for top species	64
Figure 29.	Historical comparison of Shannon diversity	66
Figure 30.	Historical comparison of species richness	67
Figure 31.	Historical comparison of fish community structure	68
Figure 32.	Relative mean abundance by fish community group	69

1.0 EXECUTIVE SUMMARY

The Vantuna Research Group at Occidental College surveyed the estuarine fishes of San Diego Bay in April and July 2019 for the Port of San Diego. The goals of the current study were to update previous studies, as well as quantify spatial and temporal utilization of the fishery populations in San Diego Bay. A summary of the results is as follows:

Composition and Abundance

During this study, 14,024 (45 species) fishes weighing a total of 271.5 kg were collected during April and July 2019. The most numerous species comprising 57.4% of the catch was Topsmelt (*Atherinops affinis*), followed by Slough Anchovy (*Anchoa delicatissima*; 9.0%), Deepbody Anchovy (*Anchoa compressa*; 7.4%), Dwarf Perch (*Micrometrus minimus*; 4.9%), and Giant Kelpfish (*Heterostichus rostratus*; 4.3%). In terms of biomass, Round Stingrays (*Urobatis halleri*) dominated the catch comprising 42.5% of the biomass, followed by Spotted Sand Bass (*Paralabrax maculatofasciatus*; 24.2%), and Topsmelt (8.4%). These species are typically among the most dominant species in surveys of San Diego Bay with the exception of Deepbody Anchovy which was only numerically dominant due to one school being captured in a single purse seine replicate.

Ecological Importance of Species

The principal fishes surveyed during these sampling periods as determined by the Ecological Index (E.I.) were the following species: Topsmelt, Round Stingray, Spotted Sand Bass, Slough Anchovy, and Deepbody Anchovy. Topsmelt ranked first (E.I. 6,572), Round Stingray ranked second (E.I. 4,606), and Spotted Sand Bass ranked third (E.I. 2,718). All three species were found ubiquitously throughout the bay during both sampling periods. Round Stingray and Spotted Sand Bass were dominant in terms of biomass, and Topsmelt was dominant in terms of numerical abundance. Slough Anchovy ranked fourth (E.I. 725) but was not present in the North Ecoregion during either sampling period. These species were followed by Deepbody Anchovy (E.I. 488), which were nearly all (98.6%) captured in a single purse seine replicate in the South-Central Ecoregion in July.

Best Estimates of Density and Standing Stock

The stock size estimate in 2019 was 18.3 million fishes, approximately half of the 2015 and 2016 estimates and the second lowest estimate ever behind 2012. With an estimated surface area of 4,858 ha this gives an overall fish density 0.38 individuals/m². The highest estimates were of Topsmelt (8.16 million), followed by Deepbody Anchovy (2.75 million), Slough Anchovy (2.19 million), Giant Kelpfish (1.09 million), and Shiner Perch (*Cymatogaster aggregata*; 0.815 million). As is typical, schooling and forage fishes dominated the stock estimate for the bay. The total best estimate of biomass standing stock was nearly 420 MT or approximately 8.64 g/m², about 35% higher than the 2016 estimate and above average for all historical surveys. The highest biomass estimates were of Round Stingray (136.5 MT), followed by Spotted Sand Bass (106.8 MT), Topsmelt (44.1 MT), Deepbody Anchovy (40.9 MT), and Bat Ray (*Myliobatis californica*; 17.0 MT). Round Stingray and Spotted Sand Bass are typically in the top five

biomass estimates from past surveys, but Deepbody Anchovy is a historically atypical top biomass species.

Avian Forage and Fisheries Species

Forage species are primarily surface-dwelling schooling fish that are accessible to diving avian predators, especially terns. Generally, forage fishes are small silvery-sided fishes that are found in large schools. These schooling fishes are not habitat-specific and move throughout the bay's ecosystem. Thirteen species of important forage fishes were captured during this study. The most abundant forage fishes were Topsmelt, Slough Anchovy, and Deepbody Anchovy. Topsmelt were primarily found at small (juvenile) size classes (<50 mm SL) appropriate for nesting birds to feed their young in the area. The typical timing for the recruitment of fishes to San Diego Bay begins in the spring and continues through the summer, but it appears this began much earlier in the year in 2019. The biomass standing stock estimate for forage fish was 106.7 MT. During this study, 13 important California recreational or commercial species were captured, the most abundant of which was Spotted Sand Bass. The standing stock estimate of fisheries species totaled 149.7 MT.

San Diego Bay as a Unique Fish Habitat and Nursery Area

San Diego Bay is known for being the northern edge of the range for many southern fishes that are not normally distributed in the Southern California Bight. Six species with primarily southern distributions were taken, including the largest Shortfin Corvina (*Cynoscion parvipinnis*; 48 cm SL) ever captured in the history of these surveys. These fishes were found almost exclusively in the southern half of the bay and none were captured in the North Ecoregion.

As the largest estuary in southern California, San Diego Bay provides critical habitat for bay and estuary fishes and continues to function as a nursery area for nearly two-thirds (65.2%) of those fishes. The high productivity rate coupled with the abundance of juvenile fishes in the bay highlights the importance of the bay as a nursery habitat. The bay contains extensive shallow water eelgrass habitat that supports a unique assemblage of juvenile and adult fishes that, in turn, support surrounding nearshore ecosystems. Juvenile fishes emigrate from the bay to offshore habitats, and important or endangered avian species utilize forage fishes in the bay. Southern California indigenous bay and estuary fishes represented 23.1% of the total catch in this survey.

Trends and Comparisons

While the 2019 surveys represent the fewest fishes ever captured in historical surveys, the biomass captured was only slightly below average. This is a product of not capturing large numbers of small individuals as is typical during April sampling, but instead capturing fewer, but larger, fish throughout the survey. This was particularly true in the South Ecoregion where fewer than 30% of the historical mean number of individuals was captured, but the biomass captured was the second highest ever recorded – more than 30% higher than average. Size structure of most species captured during the April sampling period was more typical of July sampling period catches (larger, fewer

young-of-the-year), potentially indicating a very early reproductive season for fishes in the bay.

Overall, 2019 Shannon Diversity estimates were variable among ecoregions and rank among historical values. The North and North-Central Ecoregions had the third and second lowest diversity values, respectively, of any sampling year. However, diversity values for the South-Central and South Ecoregions were the third highest of any sampling year. Species richness for 2019 was low overall (second lowest of any sampling year). Richness was the lowest ever recorded in both the North and North-Central Ecoregions and was tied for third lowest at both the South-Central and South Ecoregions for any survey period. Community structure of fishes was most similar to that of the 2012 surveys but was not statistically different than any other survey performed in the 21st Century.



Sunrise over the North Ecoregion in April 2019.

2.0 INTRODUCTION

The Vantuna Research Group at Occidental College surveyed the estuarine fishes of San Diego Bay in April and July 2019 for the Port of San Diego. The survey followed the protocols established from previous surveys (Allen 1999, Allen et al. 2002, Pondella et al. 2006, Pondella and Williams 2009a, Williams and Pondella 2012, Williams et al. 2015, Williams et al. 2016). The goals of the current study were to update the previous studies and address the following objectives:

- 1) Identify, determine, and quantify the utilization of the fishery populations in San Diego Bay
- 2) Identify habitats that support juvenile fish species and describe nursery utilization
- 3) Determine geographic and/or habitat areas of San Diego Bay that support significant populations of fish species utilized as forage by endangered avian species
- 4) Provide a comprehensive comparison of survey results to previous sampling years

In order to accomplish the objectives for these two sampling periods, we have documented the following parameters:

- ✓ Water quality parameters
- ✓ Fish species composition and abundance
 - Species diversity
 - Abundance by bay ecoregion
- ✓ Ecological importance of species
- ✓ Nursery area function
- ✓ Fish density and biomass estimates
 - Numerical and biomass density
 - Density and standing stock of avian forage species
 - Density and standing stock of fishery species
 - Panamic species unique to San Diego Bay
- ✓ Fish assemblage structure
- ✓ Historical comparisons
 - Diversity, Richness, Biomass, Abundance
 - Community structure

3.0 METHODS AND MATERIALS

3.1 Survey Locations

Four ecoregions of San Diego Bay were sampled and inventoried: North, North-Central, South-Central, and South (Figure 1, Table 1). These ecoregions were defined by Allen et al. (2002) and selected to adequately assess the status of all components of the ichthyofauna of the bay.




Figure 1. Sampling environment locations of the North, North-Central, South-Central, and South Ecoregions in San Diego Bay.

Table 1. Lambert Coordinates (Latitude, Longitude) for the San Diego Bay Fisheries Inventory and Utilization study, 2019.

Ecoregion	Site	Latitude	Longitude
North	Vegetated	32° 41' 50"	117° 13' 40"
	Non-Vegetated	32° 42' 45"	117° 12' 30"
North-Central	Vegetated	32° 41' 25"	117° 09' 50"
	Non-Vegetated	32° 41' 12"	117° 09' 45"
South-Central	Vegetated	32° 39' 05"	117° 08' 30"
	Non-Vegetated	32° 38' 48"	117° 08' 25"
South-Central	Vegetated	32° 37' 00"	117° 07' 45"
	Non-Vegetated	32° 36' 50"	117° 06' 45"

3.2 Ichthyofauna Sampling

Ichthyofauna sampling occurred during the spring and summer quarters of 2019 (April 6-7, April 13-14, and July 8-11, 2019). One ecoregion was sampled per day per sampling season. Collections were made off the 5-meter (m) *R/V Old Blue* and the 6.5-m *R/V Neoclinus*. At each ecoregion, the following five subhabitats were sampled: deep channel, nearshore non-vegetated, nearshore vegetated, intertidal non-vegetated, and intertidal vegetated.

Fish were sampled at each ecoregion using the following gear:

- 1) A 15.2-m x 1.8-m large seine equipped with a 1.8-m x 1.8-m x 1.8-m bag (1.2-centimeter [cm] mesh wings and 0.6-cm mesh in bag) was used to sample fishes in the intertidal subhabitat of each ecoregion at a depth of 0-2 meters. The net was set 15 m offshore parallel to the shoreline and pulled in shore, sampling an area of about 220 square meters (m²) per haul. Three replicates per habitat were conducted for a total of six per ecoregion.



- 2) A 4.6-m x 1.2-m small seine with 3-mm mesh was utilized to collect fish in the shallow intertidal habitat of 0-0.5m depths. The small seine was pulled 10 m along shore and pivoted towards the shore, sampling an area of approximately 62 m². Three replicates per subhabitat were conducted for a total of six per ecoregion.

- 3) A 1-m² square enclosure constructed of metal pipe and canvas was used to survey small, burrow-inhabiting fish in shallow intertidal areas of the bay. The enclosure was randomly set within each subhabitat in a depth of 0.25-0.75 m. One liter of 9:1 isopropanol-2-quinoline solution was added to the enclosed water and then searched for 10 minutes using a 1-millimeter (mm) mesh dipnet. Three replicates per subhabitat were conducted for a total of six per ecoregion.



- 4) A 1.6-m beam trawl (4-mm mesh wings and 2-mm knotless mesh in the codend) was used to sample nearshore fish species. Standardized 10-minute tows were conducted sampling an area of approximately 290 m² per replicate. Three replicates per subhabitat were conducted for a total of six per ecoregion.
- 5) A 66-m x 6-m purse seine (1.2-cm mesh wings and 0.6-cm mesh bag) was used to sample fish species in the nearshore and channel subhabitats, sampling a total area of approximately 296 m² per replicate. Three replicates per subhabitat were conducted for a total of nine per ecoregion.



- 6) An 8-m semi-balloon otter trawl (2-cm mesh wings and 0.8-cm mesh codend) was used to survey fishes from the deepest portions of the channel subhabitat. Standardized 10-minute tows were conducted sampling a total area of approximately 2,417 m² per each replicate. Three replicates were conducted per ecoregion.

All fishes were identified and measured to the nearest centimeter (standard length [SL]) and gram using measuring boards and hanging scales or a digital balance. Most individuals were measured aboard the research vessels and returned to the water, though large catches of small individuals were returned to the laboratory for identification and measurement. Coordinates of each sampling effort were recorded for all sampling events. For otter and beam trawls the start and finish of each tow were recorded. The sampling events are plotted in Figures 2-5.




Figure 2. Sampling events for the North Ecoregion, 2019.




Figure 3. Sampling events for the North-Central Ecoregion, 2019.




Figure 4. Sampling events for the South-Central Ecoregion, 2019.




Figure 5. Sampling events for the South Ecoregion, 2019.

3.3 Water Quality Sampling

Water temperature (degrees Celsius [$^{\circ}\text{C}$]), salinity (parts per thousand [ppt]), dissolved oxygen (milligrams per liter [mg/l]), and pH were measured during each sampling period and at each ecoregion using a Sea-Bird oceanographic profiler (Model SEACAT SBE 19plusV2) from *R/V Old Blue*. Downcast data from the 1-m surface bin where used to report these water quality values at the surface.

3.4 Data Analysis

3.4.1 Best Estimates of Density and Standing Stock

Density estimates used for the standing stock assessment were determined using the *Best Estimate of Density* within each Ecoregion. The best density and stock estimates were determined in the following manner:

- 1) Sample densities estimated by gear type for each species were averaged over all samples within the three depth strata (Intertidal, Nearshore, and Channel).
- 2) The maximum density for each species by gear type within each depth stratum was determined to be the *Best Estimate of Density* for that species within that depth stratum.
- 3) The proportional areal coverage of the three depth strata within the Ecoregion was determined previously by Allen et al. (2002) were used for the current study (Table 2). These areal proportions were then used to weight the *Best Estimate of Density* within the depth strata by species. A weighted average was then taken among these best estimates over the three depth strata for each species.
- 4) The sum of the weighted densities of all species represented *Best Estimate of Density* (numerical and biomass) for each depth stratum and Ecoregion was calculated.
- 5) Standing stock estimates were calculated by multiplying the best estimates by the total area of the individual Ecoregions and San Diego Bay as a whole.

Table 2. Estimates of area coverage of depth strata within each Ecoregion of San Diego Bay. Proportions and areas were used to weigh density and estimate standing stocks of fisheries.

% Area			
Ecoregion	Intertidal	Nearshore	Channel
North	6	33	60
North-Central	5	38	57
South-Central	3	61	36
South	4	84	13

Hectares/Habitat					
Ecoregion	Intertidal	Nearshore	Channel	TOTAL	% of Bay
North	61	327	593	982	20
North-Central	41	307	460	808	17
South-Central	51	1227	726	2005	41
South	40	890	133	1064	22
# Hectares	194	2751	1913	4858	
% Bay Area (Allen 2002)	4	57	39		

3.4.2 Community Structure

To characterize community-level temporal changes in fishes, we constructed a Bray-Curtis similarity matrix using the ‘vegdist’ function in the ‘vegan’ package (Oksanen et al. 2019) in R (R Core Team 2019). The matrix used fourth-root transformed taxon-specific abundance data summed across all replicates throughout the bay during each April and July. Significantly different fish community groups were determined using a cluster analysis with a SIMPROF test ($\alpha = 0.05$) performed with the ‘simprof’ function in the ‘clustsig’ package (Clarke et al. 2008; Whitaker and Christman 2014.)



Jacob Eagleton holds a 48-cm Shortfin Corvina (*Cynoscion parvipinnis*) caught by purse seine in the South Ecoregion in July 2019.

4.0 RESULTS AND DISCUSSION

4.1 Water Quality Parameters

Sea surface temperature generally increased from north to south in the bay during both sampling periods, though the temperature was about 3-5 °C warmer at each ecoregion during the July sampling period (Figure 6). Water temperature was also slightly above average in April and slightly below average in July. Dissolved oxygen was relatively consistent among Ecoregions. Salinity and pH increased slightly from north to south during the July sampling period but was consistent among ecoregions during the April sampling period. These physical-chemical results are typical of the bay and did not show effects of above-average winter rainfall.




Figure 6. Summary of physical-chemical measurements by ecoregion in April and July 2019. To contextualize the 2019 results, values from previous surveys (1995-2016) are plotted as smaller, open points.

Salinity levels in San Diego Bay are subject to the effects of seasonal rainfall and evaporation. The bay is the receiving body of water for freshwater input from three watersheds (Pueblo San Diego, Sweetwater, and Otay) covering 1,150 square kilometers (km²) of land, plus over 200 storm drain outfalls (San Diego County MS4 Co-permittees 2008). The South Ecoregion, where the Sweetwater and Otay Rivers meet the bay, has weak tidal currents, low mixing, and a small tidal prism compared to the other ecoregions (Wang et al. 1998) resulting in longer retention of freshwater signatures. However, most of the observed decreases in salinity in the South Ecoregion during the April surveys (e.g. 1998, 2012, 2015; Figure 6) were not during years where there is above average rainfall during Winter and early-Spring (e.g. 1995, 1998, 2005, 2019; Figure 7). This decoupling is due in part to the regulation of the Sweetwater and Otay Rivers by reservoirs that result in only rare releases of water into the rivers. By July, evaporation rates are typically higher than freshwater inflow creating hypersaline conditions regardless of winter precipitation levels (Peeling 1975).




Figure 7. Monthly rainfall in San Diego from 1994-2019. Data are typically collected from National Weather Service rain gauges at San Diego International Airport (Lindbergh Field). Monthly totals were compiled by John S. Stokes III.

4.2 Numerical Catch and Biomass

During this study, 14,024 (45 species) fishes weighing a total of 271.5 kilograms (kg) were collected during April and July 2019. The most numerous species comprising 57.4 percent (%) of the catch was Topsmelt (*Atherinops affinis*), followed by Slough Anchovy (*Anchoa delicatissima*; 9.0%), Deepbody Anchovy (*Anchoa compressa*; 7.4%), Dwarf Perch (*Micrometrus minimus*; 4.9%), and Giant Kelpfish (*Heterostichus rostratus*; 4.3%; Table 3). In terms of biomass, Round Stingrays (*Urobatis halleri*) dominated the catch comprising 42.5% of the biomass, followed by Spotted Sand Bass (*Paralabrax maculatofasciatus*; 24.2%), and Topsmelt (8.4%; Table 4). These species are typically among the most dominant species in surveys of San Diego Bay with the exception of Deepbody Anchovies.

Total catch decreased from north to south (Figure 8) with differences in total fish abundance reflecting the differences in Topsmelt abundance. Abundance was greatest at the North Ecoregion (6,238; Table 5), followed by the North-Central Ecoregion (4,005; Table 6), South-Central Ecoregion (2,233; Table 7), and South Ecoregion (1,548; Table 8). Topsmelt dominated catches in the North (4,867) and North-Central (2,892) Ecoregions while Deepbody Anchovy (1,019) was the dominant species in the South-Central Ecoregion where 100% of the individuals were captured in a single purse seine replicate in July. Slough Anchovy (681) was the dominant species captured in the South Ecoregion.




Figure 8. Total catch of San Diego Bay fishes by ecoregion, April and July 2019.

The catch of the five numerically dominant fishes had mixed patterns over the four ecoregions (Figure 9). Topsmelt were especially common in the northern half of the bay and Slough Anchovies were mostly captured in the southern half of the bay. Deepbody Anchovies were mostly limited to the South-Central Ecoregion where, 98.6% of all individuals caught in the entire survey were captured in a single purse seine replicate. Giant Kelpfish were found in all ecoregions and were a numerically dominant species in the North-Central Ecoregion. Dwarf Perch were limited to the North Ecoregion and were a numerically dominant species there.




Figure 9. Total catch of the five numerically dominant species by ecoregion, 2019.

Round Stingrays had the highest catch in terms of biomass at three ecoregions (North, 30.9 kg; South-Central, 19.2 kg; South, 48.8 kg) and was second in biomass (16.6 kg) to Spotted Sand Bass (25.5 kg) in the North-Central Ecoregion. Spotted Sand Bass was also a dominant species in terms of biomass in the North (6.45 kg), South-Central (12.5 kg), and South (21.2 kg) Ecoregions. Topsmelt was a dominant species in terms of biomass in the North (14.3 kg) and North-Central (7.32 kg) Ecoregions. Deepbody Anchovies (15.2 kg) and Bat Rays (*Myliobatis californica*; 8.04 kg) were among the dominant species in the South-Central and South Ecoregions, respectively.

Table 3. Total abundance of fishes collected in San Diego Bay during 2019 by ecoregion.

Scientific Name	Common Name	Ecoregions				Total	%
		North	North-Central	South-Central	South		
<i>Atherinops affinis</i>	Topsmelt	4,867	2,892	166	119	8,044	57.36
<i>Anchoa delicatissima</i>	Slough Anchovy		110	475	681	1,266	9.03
<i>Anchoa compressa</i>	Deepbody Anchovy		1	1,019	13	1,033	7.37
<i>Micrometrus minimus</i>	Dwarf Perch	691				691	4.93
<i>Heterostichus rostratus</i>	Giant Kelpfish	133	371	83	18	605	4.31
<i>Cymatogaster aggregata</i>	Shiner Perch	143	139	117	137	536	3.82
<i>Urobatis halleri</i>	Round Stingray	117	91	82	206	496	3.54
<i>Syngnathus californiensis</i>	Kelp Pipefish	82	65	121	163	431	3.07
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	37	155	81	147	420	2.99
<i>Sardinops sagax</i>	Pacific Sardine		77	1		78	0.56
<i>Paralichthys californicus</i>	California Halibut	33	12	14	5	64	0.46
<i>Clevelandia ios</i>	Arrow Goby	8	5	18	21	52	0.37
<i>Hypsoblennius gentilis</i>	Bay Blenny	20	17			37	0.26
<i>Paralabrax nebulifer</i>	Barred Sand Bass	1	27	6	1	35	0.25
<i>Leuresthes tenuis</i>	California Grunion	11	21			32	0.23
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	6	4	14	5	29	0.21
<i>Cynoscion parvipinnis</i>	Shortfin Corvina		1	18	1	20	0.14
<i>Halichoeres semicinctus</i>	Rock Wrasse	19				19	0.14
<i>Scomber japonicus</i>	Pacific Chub Mackerel	14	1			15	0.11
<i>Haemulon californiensis</i>	Salema	1	10	1	2	14	0.10
<i>Porichthys myriaster</i>	Specklefin Midshipman	11		2		13	0.09
<i>Symphurus atricaudus</i>	California Tonguefish	10	1	1		12	0.09
<i>Paralabrax clathratus</i>	Kelp Bass	9	1			10	0.07
<i>Albula gilberti</i>	Cortez Bonefish			8	1	9	0.06
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin		1	2	6	9	0.06
<i>Embiotoca jacksoni</i>	Black Perch	8				8	0.06
<i>Myliobatis californica</i>	Bat Ray			2	6	8	0.06
<i>Pleuronichthys ritteri</i>	Spotted Turbot	8				8	0.06
<i>Hippocampus ingens</i>	Pacific Seahorse				4	4	0.03
<i>Hyporhamphus rosae</i>	California Halfbeak				4	4	0.03
<i>Cheilotrema saturnum</i>	Black Croaker		2	1		3	0.02
<i>Gymnura marmorata</i>	California Butterfly Ray				3	3	0.02
<i>Fundulus parvipinnis</i>	California Killifish				2	2	0.01
<i>Scorpaena guttata</i>	California Scorpionfish	2				2	0.01
<i>Synodus lucioceps</i>	California Lizardfish	2				2	0.01
<i>Acanthogobius flavimanus</i>	Yellowfin Goby	1				1	0.01
<i>Atherinopsis californiensis</i>	Jacksmelt			1		1	0.01
<i>Dasyatis diptera</i>	Diamond Stingray				1	1	0.01
<i>Gibbonsia elegans</i>	Spotted Kelpfish	1				1	0.01
<i>Ilypnus gilberti</i>	Cheekspot Goby				1	1	0.01
<i>Phanerodon furcatus</i>	White Seaperch	1				1	0.01
<i>Pleuronichthys decurrens</i>	Curlfin Sole	1				1	0.01
<i>Seriphus politus</i>	Queenfish				1	1	0.01
<i>Umbrina roncadore</i>	Yellowfin Croaker		1			1	0.01
<i>Xystreureys liolepis</i>	Fantail Sole	1				1	0.01
# of Species: 45		6,238	4,005	2,233	1,548	14,024	

Table 4. Total biomass (grams [g]) of fishes collected in San Diego Bay during 2019 by ecoregion.

Scientific Name	Common Name	Ecoregions				Total (g)	%
		North	North-Central	South-Central	South		
<i>Urobatis halleri</i>	Round Stingray	30,909	16,586	19,194	48,775	115,464	42.53
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	6,454	25,506	12,508	21,205	65,673	24.19
<i>Atherinops affinis</i>	Topsmelt	14,319	7,320	337	716	22,691	8.36
<i>Anchoa compressa</i>	Deepbody Anchovy		10	15,200	150	15,360	5.66
<i>Myliobatis californica</i>	Bat Ray			625	8,040	8,665	3.19
<i>Albula gilberti</i>	Cortez Bonefish			4,895	490	5,385	1.98
<i>Cynoscion parvipinnis</i>	Shortfin Corvina		700	4,300	108	5,108	1.88
<i>Paralichthys californicus</i>	California Halibut	3,710	385	473	129	4,697	1.73
<i>Dasyatis diptera</i>	Diamond Stingray				4,200	4,200	1.55
<i>Micrometrus minimus</i>	Dwarf Perch	4,134				4,134	1.52
<i>Sardinops sagax</i>	Pacific Sardine		2,200	130		2,330	0.86
<i>Gymnura marmorata</i>	California Butterfly Ray				2,105	2,105	0.78
<i>Cymatogaster aggregata</i>	Shiner Perch	738	423	393	306.0	1,860	0.68
<i>Halichoeres semicinctus</i>	Rock Wrasse	1,838				1,838	0.68
<i>Anchoa delicatissima</i>	Slough Anchovy		153	703	876	1,732	0.64
<i>Heterostichus rostratus</i>	Giant Kelpfish	949	272	110	57	1,387	0.51
<i>Porichthys myriaster</i>	Specklefin Midshipman	871		505		1,376	0.51
<i>Scomber japonicus</i>	Pacific Chub Mackerel	1,200	118			1,318	0.49
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	865	293	68	58	1,284	0.47
<i>Paralabrax clathratus</i>	Kelp Bass	805	4			809	0.30
<i>Paralabrax nebulifer</i>	Barred Sand Bass	19	395	143	51	608	0.22
<i>Pleuronichthys ritteri</i>	Spotted Turbot	514				514	0.19
<i>Embiotoca jacksoni</i>	Black Perch	501				501	0.18
<i>Leuresthes tenuis</i>	California Grunion	224	198			422	0.16
<i>Syngnathus californiensis</i>	Kelp Pipefish	117	145	57	102	421	0.16
<i>Synodus lucioceps</i>	California Lizardfish	255				255	0.09
<i>Haemulon californiensis</i>	Salema	19	120	12	39	190	0.07
<i>Phanerodon furcatus</i>	White Seaperch	190				190	0.07
<i>Hypsoblennius gentilis</i>	Bay Blenny	101	46			147	0.05
<i>Scorpaena guttata</i>	California Scorpionfish	140				140	0.05
<i>Cheilotrema saturnum</i>	Black Croaker		135	< 1		135	0.05
<i>Hippocampus ingens</i>	Pacific Seahorse				134	134	0.05
<i>Symphurus atricaudus</i>	California Tonguefish	95.0	25.0	3.0		123	0.05
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin		7	19	41	67	0.02
<i>Atherinopsis californiensis</i>	Jacksmelt			66		66	0.02
<i>Umbrina roncadore</i>	Yellowfin Croaker		49			49	0.02
<i>Seriphus politus</i>	Queenfish				46	46	0.02
<i>Pleuronichthys decurrens</i>	Curlfin Sole	21				21	0.01
<i>Clevelandia ios</i>	Arrow Goby	2	3	5	9	19	0.01
<i>Hyporhamphus rosae</i>	California Halfbeak				15	15	0.01
<i>Gibbonsia elegans</i>	Spotted Kelpfish	10				10	< 0.01
<i>Xystreurus liolepis</i>	Fantail Sole	10				10	< 0.01
<i>Fundulus parvipinnis</i>	California Killifish				4	4	< 0.01
<i>Acanthogobius flavimanus</i>	Yellowfin Goby	3				3	< 0.01
<i>Ilypnus gilberti</i>	Cheekspot Goby				3	3	< 0.01
# of Species:	45	69,011	55,092	59,745	87,658	271,506	

Table 5. Total number of individuals and biomass (g) of fish species captured in the North Ecoregion, 2019.

Scientific Name	Common Name	Abundance		Biomass	
		#	%	grams	%
<i>Atherinops affinis</i>	Topsmelt	4,867	78.02	14,319	20.75
<i>Micrometrus minimus</i>	Dwarf Perch	691	11.08	4,134	5.99
<i>Cymatogaster aggregata</i>	Shiner Perch	143	2.29	738	1.07
<i>Heterostichus rostratus</i>	Giant Kelpfish	133	2.13	949	1.38
<i>Urobatis halleri</i>	Round Stingray	117	1.88	30,909	44.79
<i>Syngnathus californiensis</i>	Kelp Pipefish	82	1.31	117	0.17
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	37	0.59	6,454	9.35
<i>Paralichthys californicus</i>	California Halibut	33	0.53	3,710	5.38
<i>Hypsoblennius gentilis</i>	Bay Blenny	20	0.32	101	0.15
<i>Halichoeres semicinctus</i>	Rock Wrasse	19	0.30	1,838	2.66
<i>Scomber japonicus</i>	Pacific Chub Mackerel	14	0.22	1,200	1.74
<i>Leuresthes tenuis</i>	California Grunion	11	0.18	224	0.32
<i>Porichthys myriaster</i>	Specklefin Midshipman	11	0.18	871	1.26
<i>Symphurus atricaudus</i>	California Tonguefish	10	0.16	95	0.14
<i>Paralabrax clathratus</i>	Kelp Bass	9	0.14	805	1.17
<i>Clevelandia ios</i>	Arrow Goby	8	0.13	2	< 0.01
<i>Embiotoca jacksoni</i>	Black Perch	8	0.13	501	0.73
<i>Pleuronichthys ritteri</i>	Spotted Turbot	8	0.13	514	0.74
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	6	0.10	865	1.25
<i>Scorpaena guttata</i>	California Scorpionfish	2	0.03	140	0.20
<i>Synodus lucioceps</i>	California Lizardfish	2	0.03	255	0.37
<i>Acanthogobius flavimanus</i>	Yellowfin Goby	1	0.02	3	< 0.01
<i>Gibbonsia elegans</i>	Spotted Kelpfish	1	0.02	10	0.01
<i>Haemulon californiensis</i>	Salema	1	0.02	19	0.03
<i>Paralabrax nebulifer</i>	Barred Sand Bass	1	0.02	19	0.03
<i>Phanerodon furcatus</i>	White Seaperch	1	0.02	190	0.28
<i>Pleuronichthys decurrens</i>	Curlfin Sole	1	0.02	21	0.03
<i>Xystreurys liolepis</i>	Fantail Sole	1	0.02	10	0.01
# of Species: 28		6,238		69,011	

Table 6. Total number of individuals and biomass (g) of fish species captured in the North-Central Ecoregion, 2019.

Scientific Name	Common Name	Abundance		Biomass	
		#	%	grams	%
<i>Atherinops affinis</i>	Topsmelt	2,892	72.21	7,320	13.29
<i>Heterostichus rostratus</i>	Giant Kelpfish	371	9.26	272	0.49
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	155	3.87	25,506	46.30
<i>Cymatogaster aggregata</i>	Shiner Perch	139	3.47	423	0.77
<i>Anchoa delicatissima</i>	Slough Anchovy	110	2.75	153	0.28
<i>Urobatis halleri</i>	Round Stingray	91	2.27	16,586	30.11
<i>Sardinops sagax</i>	Pacific Sardine	77	1.92	2,200	3.99
<i>Syngnathus californiensis</i>	Kelp Pipefish	65	1.62	145	0.26
<i>Paralabrax nebulifer</i>	Barred Sand Bass	27	0.67	395	0.72
<i>Leuresthes tenuis</i>	California Grunion	21	0.52	198	0.36
<i>Hypsoblennius gentilis</i>	Bay Blenny	17	0.42	46	0.08
<i>Paralichthys californicus</i>	California Halibut	12	0.30	385	0.70
<i>Haemulon californiensis</i>	Salema	10	0.25	120	0.22
<i>Clevelandia ios</i>	Arrow Goby	5	0.12	3	0.01
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	4	0.10	293	0.53
<i>Cheilotrema saturnum</i>	Black Croaker	2	0.05	135	0.25
<i>Anchoa compressa</i>	Deepbody Anchovy	1	0.02	10	0.02
<i>Cynoscion parvipinnis</i>	Shortfin Corvina	1	0.02	700	1.27
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin	1	0.02	7	0.01
<i>Paralabrax clathratus</i>	Kelp Bass	1	0.02	4	0.01
<i>Scomber japonicus</i>	Pacific Chub Mackerel	1	0.02	118	0.21
<i>Symphurus atricaudus</i>	California Tonguefish	1	0.02	25	0.05
<i>Umbrina roncadior</i>	Yellowfin Croaker	1	0.02	49	0.09
# of Species: 23		4,005		55,092	

Table 7. Total number of individuals and biomass (g) of fish species captured in the South-Central Ecoregion, 2019.

Scientific Name	Common Name	Abundance		Biomass	
		#	%	grams	%
<i>Anchoa compressa</i>	Deepbody Anchovy	1,019	45.63	15,200	25.44
<i>Anchoa delicatissima</i>	Slough Anchovy	475	21.27	703	1.18
<i>Atherinops affinis</i>	Topsmelt	166	7.43	337	0.56
<i>Syngnathus californiensis</i>	Kelp Pipefish	121	5.42	57	0.10
<i>Cymatogaster aggregata</i>	Shiner Perch	117	5.24	393	0.66
<i>Heterostichus rostratus</i>	Giant Kelpfish	83	3.72	110	0.18
<i>Urobatis halleri</i>	Round Stingray	82	3.67	19,194	32.13
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	81	3.63	12,508	20.94
<i>Clevelandia ios</i>	Arrow Goby	18	0.81	5	0.01
<i>Cynoscion parvipinnis</i>	Shortfin Corvina	18	0.81	4,300	7.20
<i>Paralichthys californicus</i>	California Halibut	14	0.63	473	0.79
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	14	0.63	68	0.11
<i>Albula gilberti</i>	Cortez Bonefish	8	0.36	4,895	8.19
<i>Paralabrax nebulifer</i>	Barred Sand Bass	6	0.27	143	0.24
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin	2	0.09	19	0.03
<i>Myliobatis californica</i>	Bat Ray	2	0.09	625	1.05
<i>Porichthys myriaster</i>	Specklefin Midshipman	2	0.09	505	0.85
<i>Atherinopsis californiensis</i>	Jacksmelt	1	0.04	66	0.11
<i>Cheilotrema saturnum</i>	Black Croaker	1	0.04	0.1	< 0.01
<i>Haemulon californiensis</i>	Salema	1	0.04	12	0.02
<i>Sardinops sagax</i>	Pacific Sardine	1	0.04	130	0.22
<i>Symphurus atricaudus</i>	California Tonguefish	1	0.04	3	0.01
# of Species: 22		2,233		59,745	

Table 8. Total number of individuals and biomass (g) of fish species captured in the South Ecoregion, 2019.

Scientific Name	Common Name	Abundance		Biomass	
		#	%	grams	%
<i>Anchoa delicatissima</i>	Slough Anchovy	681	43.99	876	1.00
<i>Urobatis halleri</i>	Round Stingray	206	13.31	48,775	55.64
<i>Syngnathus californiensis</i>	Kelp Pipefish	163	10.53	102	0.12
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	147	9.50	21,205	24.19
<i>Cymatogaster aggregata</i>	Shiner Perch	137	8.85	306	0.35
<i>Atherinops affinis</i>	Topsmelt	119	7.69	716	0.82
<i>Clevelandia ios</i>	Arrow Goby	21	1.36	9	0.01
<i>Heterostichus rostratus</i>	Giant Kelpfish	18	1.16	57	0.07
<i>Anchoa compressa</i>	Deepbody Anchovy	13	0.84	150	0.17
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin	6	0.39	41	0.05
<i>Myliobatis californica</i>	Bat Ray	6	0.39	8,040	9.17
<i>Paralichthys californicus</i>	California Halibut	5	0.32	129	0.15
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	5	0.32	58	0.07
<i>Hippocampus ingens</i>	Pacific Seahorse	4	0.26	134	0.15
<i>Hyporhamphus rosae</i>	California Halfbeak	4	0.26	15	0.02
<i>Gymnura marmorata</i>	California Butterfly Ray	3	0.19	2,105	2.40
<i>Fundulus parvipinnis</i>	California Killifish	2	0.13	4	< 0.01
<i>Haemulon californiensis</i>	Salema	2	0.13	39	0.04
<i>Albula gilberti</i>	Cortez Bonefish	1	0.06	490	0.56
<i>Cynoscion parvipinnis</i>	Shortfin Corvina	1	0.06	108	0.12
<i>Dasyatis dipterura</i>	Diamond Stingray	1	0.06	4,200	4.79
<i>Ilypnus gilberti</i>	Cheekspot Goby	1	0.06	3	< 0.01
<i>Paralabrax nebulifer</i>	Barred Sand Bass	1	0.06	51	0.06
<i>Seriphus politus</i>	Queenfish	1	0.06	46	0.05
# of Species: 24		1,548		87,658	

4.3 Shannon Diversity and Species Richness

The Shannon Diversity index (H' ; Shannon 1948) was used to estimate diversity in San Diego Bay and provide a basis for comparison among ecoregions within the bay. Diversity was calculated for total catches by ecoregion and by sampling month. Species richness was low overall in 2019 but followed a typical spatial pattern with richness decreasing from north to south. Diversity, however, was lowest in the North Ecoregion and increased to the south (Figure 10). Species richness decreased slightly while H' increased from April to July 2019 (Figure 11). Both temporally and spatially, reduced H' values were due to the dominance of forage fishes.




Figure 10. Shannon Diversity (H') and number of species (richness) of fishes taken in each San Diego Bay ecoregion, 2019.




Figure 11. Shannon Diversity (H') and number of species (richness) of fishes taken in San Diego Bay by sampling month, 2019.

4.4 Catch by Sampling Ecoregion and Period

North Ecoregion – A total of 4,867 fishes belonging to 28 species and weighing 69.0 kg was collected in the North Ecoregion over two sampling periods in 2019 (Table 4). Topsmelt was by far the most frequently caught species (78.0%), followed by Dwarf Perch (11.1%), Shiner Perch (*Cymatogaster aggregata*; 2.29%), Giant Kelpfish (2.13%), and Round Stingray (1.88%). Round Stingray led in total biomass (44.8%), followed by Topsmelt (20.8%), Spotted Sand Bass (9.35%), Dwarf Perch (5.99%) and California Halibut (*Paralichthys californicus*; 5.38%)

North-Central Ecoregion - A total of 4,005 fishes belonging to 23 species and weighing 55.1 kg was collected in the North-Central Ecoregion in April and July, 2019 (Table 5). Topsmelt was by far the most abundant species (72.2%), followed by Giant Kelpfish (9.26%), Spotted Sand Bass (3.87%), Shiner Perch (3.47%), and Slough Anchovy (2.75%). Spotted Sand Bass led in total biomass (46.3%), followed by Round Stingray (30.1%), Topsmelt (13.3%), and Pacific Sardine (*Sardinops sagax*; 3.99%).

South-Central Ecoregion - A total of 2,233 fishes belonging to 22 species and weighing 59.7 kg was collected in the South-Central Ecoregion over the two sampling periods in 2019 (Table 6). Deepbody Anchovy was the most abundant species (45.6%), followed by Slough Anchovy (21.3%), Topsmelt (7.43%), Kelp Pipefish (*Syngnathus californiensis*; 5.42%), and Shiner Perch (5.24%). Round Stingray led in total biomass (32.1%), followed by Deepbody Anchovy (25.4%), Spotted Sand Bass (20.9%), Cortez Bonefish (*Albula gilberti*; 8.19%), and Shortfin Corvina (*Cynoscion parvipinnis*; 7.20%).

South Ecoregion - A total of 1,548 fishes belonging to 24 species and weighing 87.7 kg was collected in the South Ecoregion in April and July 2019 (Table 7). Slough Anchovy was the most abundant species (44.0%), followed Round Stingray (13.3%), Kelp Pipefish (10.5%), Spotted Sand Bass (9.50%), and Shiner Perch (8.85%). Round Stingray led in total biomass (55.6%), followed by Spotted Sand Bass (24.2%), and Bat Ray (9.2%).

In April 2019, 8,159 individuals comprised of 37 species of fishes were captured (Figure 12, Table 9). In July, the catch decreased to 5,865 fish and species richness decreased slightly to 35. Total biomass was greater in April (141.7 kg) than July (129.8 kg) (Figure 12, Table 10). From April to July there was a substantial decrease in total biomass in the North Ecoregion (Figure 13) presumably due to Topsmelt moving from the warm, protected waters towards the open ocean over this period. Biomass remained relatively steady at the other three ecoregions.




Figure 12. Total catch of fishes and biomass (kg) taken in San Diego Bay by sampling period, 2019.




Figure 13. Biomass (kg) of San Diego Bay fishes taken by ecoregion, April and July 2019.

Table 9. Total abundance of fish species taken in San Diego Bay by sampling period, 2019.

Scientific Name	Common Name	2019		Total	%
		April	July		
<i>Atherinops affinis</i>	Topsmelt	5,357	2,687	8,044	57.36
<i>Anchoa delicatissima</i>	Slough Anchovy	641	625	1,266	9.03
<i>Anchoa compressa</i>	Deepbody Anchovy	1	1,032	1,033	7.37
<i>Micrometrus minimus</i>	Dwarf Perch	136	555	691	4.93
<i>Heterostichus rostratus</i>	Giant Kelpfish	497	108	605	4.31
<i>Cymatogaster aggregata</i>	Shiner Perch	445	91	536	3.82
<i>Urobatis halleri</i>	Round Stingray	254	242	496	3.54
<i>Syngnathus californiensis</i>	Kelp Pipefish	264	167	431	3.07
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	233	187	420	2.99
<i>Sardinops sagax</i>	Pacific Sardine	78		78	0.56
<i>Paralichthys californicus</i>	California Halibut	32	32	64	0.46
<i>Clevelandia ios</i>	Arrow Goby	23	29	52	0.37
<i>Hypsoblennius gentilis</i>	Bay Blenny	31	6	37	0.26
<i>Paralabrax nebulifer</i>	Barred Sand Bass	24	11	35	0.25
<i>Leuresthes tenuis</i>	California Grunion	32		32	0.23
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	21	8	29	0.21
<i>Cynoscion parvipinnis</i>	Shortfin Corvina	1	19	20	0.14
<i>Halichoeres semicinctus</i>	Rock Wrasse	7	12	19	0.14
<i>Scomber japonicus</i>	Pacific Chub Mackerel	14	1	15	0.11
<i>Haemulon californiensis</i>	Salema	12	2	14	0.10
<i>Porichthys myriaster</i>	Specklefin Midshipman	4	9	13	0.09
<i>Symphurus atricaudus</i>	California Tonguefish	2	10	12	0.09
<i>Paralabrax clathratus</i>	Kelp Bass	7	3	10	0.07
<i>Albula gilberti</i>	Cortez Bonefish	6	3	9	0.06
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin	8	1	9	0.06
<i>Embiotoca jacksoni</i>	Black Perch	3	5	8	0.06
<i>Myliobatis californica</i>	Bat Ray	6	2	8	0.06
<i>Pleuronichthys ritteri</i>	Spotted Turbot	3	5	8	0.06
<i>Hippocampus ingens</i>	Pacific Seahorse	4		4	0.03
<i>Hyporhamphus rosae</i>	California Halfbeak	4		4	0.03
<i>Cheilotrema saturnum</i>	Black Croaker		3	3	0.02
<i>Gymnura marmorata</i>	California Butterfly Ray	3		3	0.02
<i>Fundulus parvipinnis</i>	California Killifish		2	2	0.01
<i>Scorpaena guttata</i>	California Scorpionfish	1	1	2	0.01
<i>Synodus lucioceps</i>	California Lizardfish		2	2	0.01
<i>Acanthogobius flavimanus</i>	Yellowfin Goby	1		1	0.01
<i>Atherinopsis californiensis</i>	Jacksmelt		1	1	0.01
<i>Dasyatis dipterura</i>	Diamond Stingray	1		1	0.01
<i>Gibbonsia elegans</i>	Spotted Kelpfish	1		1	0.01
<i>Ilypnus gilberti</i>	Cheekspot Goby		1	1	0.01
<i>Phanerodon furcatus</i>	White Seaperch	1		1	0.01
<i>Pleuronichthys decurrens</i>	Curlfin Sole		1	1	0.01
<i>Seriphus politus</i>	Queenfish	1		1	0.01
<i>Umbrina roncadior</i>	Yellowfin Croaker		1	1	0.01
<i>Xystreurus liolepis</i>	Fantail Sole		1	1	0.01
Total:		8,159	5,865	14,024	
# of Species:		37	35		

Table 10. Total biomass (g) of fish species taken in San Diego Bay by sampling period, 2019.

Scientific Name	Common Name	2019		Total	%
		April	July		
<i>Urobatis halleri</i>	Round Stingray	50,731	64,733	115,464	42.53
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	35,308	30,365	65,673	24.19
<i>Atherinops affinis</i>	Topsmelt	20,396	2,295	22,691	8.36
<i>Anchoa compressa</i>	Deepbody Anchovy	10	15,350	15,360	5.66
<i>Myliobatis californica</i>	Bat Ray	8,040	625	8,665	3.19
<i>Albula gilberti</i>	Cortez Bonefish	3,540	1,845	5,385	1.98
<i>Cynoscion parvipinnis</i>	Shortfin Corvina	700	4,408	5,108	1.88
<i>Paralichthys californicus</i>	California Halibut	3,431	1,266	4,697	1.73
<i>Dasyatis diptera</i>	Diamond Stingray	4,200		4,200	1.55
<i>Micrometrus minimus</i>	Dwarf Perch	1,772	2,362	4,134	1.52
<i>Sardinops sagax</i>	Pacific Sardine	2,330		2,330	0.86
<i>Gymnura marmorata</i>	California Butterfly Ray	2,105		2,105	0.78
<i>Cymatogaster aggregata</i>	Shiner Perch	1,290	570	1,860	0.68
<i>Halichoeres semicinctus</i>	Rock Wrasse	239	1,599	1,838	0.68
<i>Anchoa delicatissima</i>	Slough Anchovy	792	940	1,732	0.64
<i>Heterostichus rostratus</i>	Giant Kelpfish	665	722	1,387	0.51
<i>Porichthys myriaster</i>	Specklefin Midshipman	1,237	139	1,376	0.51
<i>Scomber japonicus</i>	Pacific Chub Mackerel	1,200	118	1,318	0.49
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	708	576	1,284	0.47
<i>Paralabrax clathratus</i>	Kelp Bass	742	67	809	0.30
<i>Paralabrax nebulifer</i>	Barred Sand Bass	302	306	608	0.22
<i>Pleuronichthys ritteri</i>	Spotted Turbot	265	249	514	0.19
<i>Embiotoca jacksoni</i>	Black Perch	328	173	501	0.18
<i>Leuresthes tenuis</i>	California Grunion	422		422	0.16
<i>Syngnathus californiensis</i>	Kelp Pipefish	247	175	421	0.16
<i>Synodus lucioceps</i>	California Lizardfish		255	255	0.09
<i>Haemulon californiensis</i>	Salema	151	39	190	0.07
<i>Phanerodon furcatus</i>	White Seaperch	190		190	0.07
<i>Hypsoblennius gentilis</i>	Bay Blenny	52	95	147	0.05
<i>Scorpaena guttata</i>	California Scorpionfish	5	135	140	0.05
<i>Cheilotrema saturnum</i>	Black Croaker		135	135	0.05
<i>Hippocampus ingens</i>	Pacific Seahorse	134		134	0.05
<i>Symphurus atricaudus</i>	California Tonguefish	40	83	123	0.05
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin	60	7	67	0.02
<i>Atherinopsis californiensis</i>	Jacksnelt		66	66	0.02
<i>Umbrina roncadore</i>	Yellowfin Croaker		49	49	0.02
<i>Seriphus politus</i>	Queenfish	46		46	0.02
<i>Pleuronichthys decurrens</i>	Curlfin Sole		21	21	0.01
<i>Clevelandia ios</i>	Arrow Goby	8	11	19	0.01
<i>Hyporhamphus rosae</i>	California Halfbeak	15		15	0.01
<i>Gibbonsia elegans</i>	Spotted Kelpfish	10		10	< 0.01
<i>Xystreureys liolepis</i>	Fantail Sole		10	10	< 0.01
<i>Fundulus parvipinnis</i>	California Killifish		4	4	< 0.01
<i>Acanthogobius flavimanus</i>	Yellowfin Goby	3		3	< 0.01
<i>Ilypnus gilberti</i>	Cheekspot Goby		3	3	< 0.01
Total:		141,713	129,794	271,506	
# of Species:		37	35		

4.5 Catch by Depth Strata and Subhabitats

Of the three bay depth strata (intertidal, nearshore, and channel) the greatest catch of fishes was in the intertidal strata (6,315 individuals from 24 species; Table 11). 5,876 fishes from 33 species were captured in the nearshore, and 1,833 fishes from 24 species were captured in the channel. A total of 8,076 fishes was taken in vegetated areas of the nearshore and intertidal (Table 12) comprised of 28 of the 45 species captured during the 2019 surveys. 4,115 fishes, from 33 species were caught in the nearshore and intertidal non-vegetated areas.



Purse seine being pulled aboard R/V Neoclinus from the channel in the South-Central Ecoregion in July 2019.

Table 11. Total abundance of fish species taken from San Diego Bay by depth strata, 2019.

Scientific Name	Common Name	Depth Strata			Total	%
		Channel	Intertidal	Nearshore		
<i>Atherinops affinis</i>	Topsmelt	278	5,480	2,286	8,044	57.36
<i>Anchoa delicatissima</i>	Slough Anchovy	124	206	936	1,266	9.03
<i>Anchoa compressa</i>	Deepbody Anchovy	1,032	1		1,033	7.37
<i>Micrometrus minimus</i>	Dwarf Perch		334	357	691	4.93
<i>Heterostichus rostratus</i>	Giant Kelpfish		39	566	605	4.31
<i>Cymatogaster aggregata</i>	Shiner Perch	4	11	521	536	3.82
<i>Urobatis halleri</i>	Round Stingray	159	12	325	496	3.54
<i>Syngnathus californiensis</i>	Kelp Pipefish	1	115	315	431	3.07
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	59	5	356	420	2.99
<i>Sardinops sagax</i>	Pacific Sardine	78			78	0.56
<i>Paralichthys californicus</i>	California Halibut	25	14	25	64	0.46
<i>Clevelandia ios</i>	Arrow Goby		38	14	52	0.37
<i>Hypsoblennius gentilis</i>	Bay Blenny	1	5	31	37	0.26
<i>Paralabrax nebulifer</i>	Barred Sand Bass		9	26	35	0.25
<i>Leuresthes tenuis</i>	California Grunion		3	29	32	0.23
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	10	18	1	29	0.21
<i>Cynoscion parvipinnis</i>	Shortfin Corvina	17		3	20	0.14
<i>Halichoeres semicinctus</i>	Rock Wrasse			19	19	0.14
<i>Scomber japonicus</i>	Pacific Chub Mackerel	15			15	0.11
<i>Haemulon californiensis</i>	Salema			14	14	0.10
<i>Porichthys myriaster</i>	Specklefin Midshipman	3		10	13	0.09
<i>Symphurus atricaudus</i>	California Tonguefish	2		10	12	0.09
<i>Paralabrax clathratus</i>	Kelp Bass			10	10	0.07
<i>Albula gilberti</i>	Cortez Bonefish	7		2	9	0.06
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin		9		9	0.06
<i>Embiotoca jacksoni</i>	Black Perch		5	3	8	0.06
<i>Myliobatis californica</i>	Bat Ray	2	1	5	8	0.06
<i>Pleuronichthys ritteri</i>	Spotted Turbot	7		1	8	0.06
<i>Hippocampus ingens</i>	Pacific Seahorse	1		3	4	0.03
<i>Hyporhamphus rosae</i>	California Halfbeak		4		4	0.03
<i>Cheilotrema saturnum</i>	Black Croaker	2		1	3	0.02
<i>Gymnura marmorata</i>	California Butterfly Ray	1	1	1	3	0.02
<i>Fundulus parvipinnis</i>	California Killifish		2		2	0.01
<i>Scorpaena guttata</i>	California Scorpionfish	2			2	0.01
<i>Synodus lucioceps</i>	California Lizardfish	2			2	0.01
<i>Acanthogobius flavimanus</i>	Yellowfin Goby			1	1	0.01
<i>Atherinopsis californiensis</i>	Jacksmelt		1		1	0.01
<i>Dasyatis dipterura</i>	Diamond Stingray			1	1	0.01
<i>Gibbonsia elegans</i>	Spotted Kelpfish			1	1	0.01
<i>Ilypnus gilberti</i>	Cheekspot Goby		1		1	0.01
<i>Phanerodon furcatus</i>	White Seaperch			1	1	0.01
<i>Pleuronichthys decurrens</i>	Curlfin Sole			1	1	0.01
<i>Seriphus politus</i>	Queenfish			1	1	0.01
<i>Umbrina roncadore</i>	Yellowfin Croaker		1		1	0.01
<i>Xystreureys liolepis</i>	Fantail Sole	1			1	0.01
Total:		1,833	6,315	5,876	14,024	
# of Species:		24	24	33		

Table 12. Total abundance of fish species taken from San Diego Bay by subhabitat, 2019.


Scientific Name	Common Name	Subhabitat			Total	%
		Channel	Non-Vegetated	Vegetated		
<i>Atherinops affinis</i>	Topsmelt	278	1,986	5,780	8,044	57.36
<i>Anchoa delicatissima</i>	Slough Anchovy	124	730	412	1,266	9.03
<i>Anchoa compressa</i>	Deepbody Anchovy	1,032		1	1,033	7.37
<i>Micrometrus minimus</i>	Dwarf Perch		254	437	691	4.93
<i>Heterostichus rostratus</i>	Giant Kelpfish		293	312	605	4.31
<i>Cymatogaster aggregata</i>	Shiner Perch	4	253	279	536	3.82
<i>Urobatis halleri</i>	Round Stingray	159	69	268	496	3.54
<i>Syngnathus californiensis</i>	Kelp Pipefish	1	210	220	431	3.07
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	59	143	218	420	2.99
<i>Sardinops sagax</i>	Pacific Sardine	78			78	0.56
<i>Paralichthys californicus</i>	California Halibut	25	27	12	64	0.46
<i>Clevelandia ios</i>	Arrow Goby		21	31	52	0.37
<i>Hypsoblennius gentilis</i>	Bay Blenny	1	25	11	37	0.26
<i>Paralabrax nebulifer</i>	Barred Sand Bass		19	16	35	0.25
<i>Leuresthes tenuis</i>	California Grunion		6	26	32	0.23
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	10	5	14	29	0.21
<i>Cynoscion parvipinnis</i>	Shortfin Corvina	17	1	2	20	0.14
<i>Halichoeres semicinctus</i>	Rock Wrasse		12	7	19	0.14
<i>Scomber japonicus</i>	Pacific Chub Mackerel	15			15	0.11
<i>Haemulon californiensis</i>	Salema		10	4	14	0.10
<i>Porichthys myriaster</i>	Specklefin Midshipman	3	10		13	0.09
<i>Symphurus atricaudus</i>	California Tonguefish	2	7	3	12	0.09
<i>Paralabrax clathratus</i>	Kelp Bass		4	6	10	0.07
<i>Albula gilberti</i>	Cortez Bonefish	7	2		9	0.06
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin		4	5	9	0.06
<i>Embiotoca jacksoni</i>	Black Perch		4	4	8	0.06
<i>Myliobatis californica</i>	Bat Ray	2	6		8	0.06
<i>Pleuronichthys ritteri</i>	Spotted Turbot	7	1		8	0.06
<i>Hippocampus ingens</i>	Pacific Seahorse	1	3		4	0.03
<i>Hyporhamphus rosae</i>	California Halfbeak		4		4	0.03
<i>Cheilotrema saturnum</i>	Black Croaker	2	1		3	0.02
<i>Gymnura marmorata</i>	California Butterfly Ray	1		2	3	0.02
<i>Fundulus parvipinnis</i>	California Killifish			2	2	0.01
<i>Scorpaena guttata</i>	California Scorpionfish	2			2	0.01
<i>Synodus lucioceps</i>	California Lizardfish	2			2	0.01
<i>Acanthogobius flavimanus</i>	Yellowfin Goby		1		1	0.01
<i>Atherinopsis californiensis</i>	Jacksmelt			1	1	0.01
<i>Dasyatis dipterura</i>	Diamond Stingray		1		1	0.01
<i>Gibbonsia elegans</i>	Spotted Kelpfish			1	1	0.01
<i>Ilypnus gilberti</i>	Cheekspot Goby			1	1	0.01
<i>Phanerodon furcatus</i>	White Seaperch			1	1	0.01
<i>Pleuronichthys decurrens</i>	Curlfin Sole		1		1	0.01
<i>Seriphus politus</i>	Queenfish		1		1	0.01
<i>Umbrina roncadore</i>	Yellowfin Croaker		1		1	0.01
<i>Xystreureys liolepis</i>	Fantail Sole	1			1	0.01
Total:		1,833	4,115	8,076	14,024	
# of Species:		24	33	28		

4.6 Nursery Area Function

San Diego Bay continues to be a nursery area for over half of the fishes found there. Approximately 65% of all fishes sampled in San Diego Bay were juveniles (Table 13). The most abundant juveniles (Topsmelt, Arrow Goby, Giant Kelpfish) are all critical commercial and/or forage fish species. The high catch of juvenile fishes in the bay highlights the continued importance of San Diego Bay as a nursery area for bay, estuarine, and nearshore species.

Table 13. Percent of juveniles taken of the top 20 species of fish from San Diego Bay, 2019.

Scientific Name	Common Name	Total		
		Juveniles	Abundance	% Juvenile
<i>Hypsoblennius gentilis</i>	Bay Blenny	37	37	100.00
<i>Clevelandia ios</i>	Arrow Goby	49	52	94.23
<i>Heterostichus rostratus</i>	Giant Kelpfish	569	605	94.05
<i>Haemulon californiensis</i>	Salema	13	14	92.86
<i>Atherinops affinis</i>	Topsmelt	7,166	8,044	89.09
<i>Paralabrax nebulifer</i>	Barred Sand Bass	30	35	85.71
<i>Cymatogaster aggregata</i>	Shiner Perch	414	536	77.24
<i>Micrometrus minimus</i>	Dwarf Perch	502	691	72.65
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	21	29	72.41
<i>Paralichthys californicus</i>	California Halibut	36	64	56.25
<i>Syngnathus californiensis</i>	Kelp Pipefish	204	431	47.33
<i>Halichoeres semicinctus</i>	Rock Wrasse	6	19	31.58
<i>Urobatis halleri</i>	Round Stingray	21	496	4.23
<i>Leuresthes tenuis</i>	California Grunion	1	32	3.13
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	7	420	1.67
<i>Anchoa delicatissima</i>	Slough Anchovy	2	1,266	0.16
<i>Anchoa compressa</i>	Deepbody Anchovy	0	1,033	0.00
<i>Sardinops sagax</i>	Pacific Sardine	0	78	0.00
<i>Cynoscion parvipinnis</i>	Shortfin Corvina	0	20	0.00
<i>Scomber japonicus</i>	Pacific Chub Mackerel	0	15	0.00
		9,078	13,917	65.23



Juvenile Giant Kelpfish captured by large seine from the South-Central Ecoregion in April 2019.

4.7 Ecological Importance of Species

An index of ecological importance was also calculated to estimate the relative importance of each species within the bay assemblage. An Ecological Index (E.I.) was determined using the total catch for each species during this study and incorporated three significant ecological variables: % Number, % Weight, and % Frequency of Occurrence, by ecoregion and month ($E.I. = (\% \text{ Number} + \% \text{ Weight}) * \% \text{ Frequency of Occurrence}$; Table 14; Figure 14). This index is indicative of the importance of each species to the energy flow within the San Diego Bay ecosystem. Topsmelt ranked first (E.I. 6,572), Round Stingray ranked second (E.I. 4,606), and Spotted Sand Bass ranked third (E.I. 2,718). All three species were found ubiquitously throughout the bay during both sampling periods. Round Stingray and Spotted Sand Bass were dominant in terms of biomass, and Topsmelt was dominant in terms of numerical abundance. Slough Anchovy ranked fourth (E.I. 725) but was not present in the North Ecoregion during either sampling period. These species were followed by Deepbody Anchovy (E.I. 488), which were nearly all (98.6%) captured in a single purse seine replicate in the South-Central Ecoregion in July.




Figure 14. Top 20 species of San Diego Bay fishes ranked by Ecological Index, 2019.

Table 14. Relative abundance, relative biomass, frequency of occurrence, and Ecological Index (E.I.) of San Diego Bay fishes, 2019.

Scientific Name	Common Name	Abundance %	Biomass %	Frequency of Occurrence	Ecological Index
<i>Atherinops affinis</i>	Topsmelt	57.36	8.36	100.0	6571.61
<i>Urobatis halleri</i>	Round Stingray	3.54	42.53	100.0	4606.40
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	2.99	24.19	100.0	2718.31
<i>Anchoa delicatissima</i>	Slough Anchovy	9.03	0.64	75.0	724.90
<i>Anchoa compressa</i>	Deepbody Anchovy	7.37	5.66	37.5	488.37
<i>Heterostichus rostratus</i>	Giant Kelpfish	4.31	0.51	100.0	482.49
<i>Cymatogaster aggregata</i>	Shiner Perch	3.82	0.68	87.5	394.35
<i>Syngnathus californiensis</i>	Kelp Pipefish	3.07	0.16	100.0	322.84
<i>Paralichthys californicus</i>	California Halibut	0.46	1.73	87.5	191.29
<i>Micrometrus minimus</i>	Dwarf Perch	4.93	1.52	25.0	161.25
<i>Myliobatis californica</i>	Bat Ray	0.06	3.19	25.0	81.21
<i>Albula gilberti</i>	Cortez Bonefish	0.06	1.98	37.5	76.78
<i>Cynoscion parvipinnis</i>	Shortfin Corvina	0.14	1.88	37.5	75.90
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	0.21	0.47	87.5	59.47
<i>Paralabrax nebulifer</i>	Barred Sand Bass	0.25	0.22	75.0	35.51
<i>Sardinops sagax</i>	Pacific Sardine	0.56	0.86	25.0	35.36
<i>Clevelandia ios</i>	Arrow Goby	0.37	0.01	75.0	28.32
<i>Porichthys myriaster</i>	Specklefin Midshipman	0.09	0.51	37.5	22.47
<i>Halichoeres semicinctus</i>	Rock Wrasse	0.14	0.68	25.0	20.31
<i>Dasyatis dipterura</i>	Diamond Stingray	0.01	1.55	12.5	19.43
<i>Hypsoblennius gentilis</i>	Bay Blenny	0.26	0.05	50.0	15.90
<i>Scomber japonicus</i>	Pacific Chub Mackerel	0.11	0.49	25.0	14.81
<i>Paralabrax clathratus</i>	Kelp Bass	0.07	0.30	37.5	13.85
<i>Gymnura marmorata</i>	California Butterfly Ray	0.02	0.78	12.5	9.96
<i>Leuresthes tenuis</i>	California Grunion	0.23	0.16	25.0	9.59
<i>Haemulon californiensis</i>	Salema	0.10	0.07	50.0	8.49
<i>Symphurus atricaudus</i>	California Tonguefish	0.09	0.05	50.0	6.54
<i>Pleuronichthys ritteri</i>	Spotted Turbot	0.06	0.19	25.0	6.16
<i>Embiotoca jacksoni</i>	Black Perch	0.06	0.18	25.0	6.04
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin	0.06	0.02	37.5	3.33
<i>Cheilotrema saturnum</i>	Black Croaker	0.02	0.05	25.0	1.78
<i>Scorpaena guttata</i>	California Scorpionfish	0.01	0.05	25.0	1.65
<i>Synodus lucioceps</i>	California Lizardfish	0.01	0.09	12.5	1.35
<i>Hippocampus ingens</i>	Pacific Seahorse	0.03	0.05	12.5	0.97
<i>Phanerodon furcatus</i>	White Seaperch	0.01	0.07	12.5	0.96
<i>Hyporhamphus rosae</i>	California Halfbeak	0.03	0.01	12.5	0.43
<i>Atherinopsis californiensis</i>	Jacksmelt	0.01	0.02	12.5	0.39
<i>Umbrina roncadore</i>	Yellowfin Croaker	0.01	0.02	12.5	0.31
<i>Seriphus politus</i>	Queenfish	0.01	0.02	12.5	0.30
<i>Fundulus parvipinnis</i>	California Killifish	0.01	< 0.01	12.5	0.20
<i>Pleuronichthys decurrens</i>	Curffin Sole	0.01	0.01	12.5	0.19
<i>Gibbonsia elegans</i>	Spotted Kelpfish	0.01	< 0.01	12.5	0.14
<i>Xystreureys liolepis</i>	Fantail Sole	0.01	< 0.01	12.5	0.14
<i>Acanthogobius flavimanus</i>	Yellowfin Goby	0.01	< 0.01	12.5	0.10
<i>Ilypnus gilberti</i>	Cheekspot Goby	0.01	< 0.01	12.5	0.10

4.8 Principal species

Topsmelt (*Atherinops affinis*)

Topsmelt was ranked first by the Ecological Index and were found during both sampling periods and at all ecoregions, depth strata and subhabitats, though the vast majority were caught in the intertidal depth strata. These mostly juvenile



fish (89.1%) were nearly three times as abundant in the vegetated (5,780) versus non-vegetated (1,986) subhabitats. They were the most abundant fish in the survey comprising 57.4% of the catch, but only 8.36% of the biomass due to their relatively small size. Their size frequency distribution (Figure 15) in April showed a pattern typically more consistent with that of July, suggesting that the reproductive season for Topsmelt occurred earlier than normal in 2019. No adults were observed in July.




Figure 15. Total number of Topsmelt individuals by standard length (cm) from San Diego Bay, April and July 2019.

Round Stingray (*Urobatis halleri*)



The second highest ranked species in terms of ecological importance in 2019 was the Round Stingray. This species was ubiquitous throughout the bay during these surveys, and were found in all sampling periods, ecoregions, depth strata and subhabitats. While only consisting of 3.54% of the total individuals captured in the 2019 surveys, those individuals accounted for 42.5% of the biomass. The sizes of

captured Round Stingrays were widely stratified, representative of its entire size range, and showed growth of juveniles and sub-adults between the April and July sampling periods (Figure 16). Round Stingrays were caught primarily in the channel and nearshore depth strata, with just a few individuals captured in the intertidal. While they were observed in all four ecoregions, the highest catches were in the South Ecoregion.





Figure 16. Total number of Round Stingray individuals by disc width (cm) from San Diego Bay, April and July 2019.

Spotted Sand Bass (*Paralabrax maculatofasciatus*)

Spotted Sand Bass are the ubiquitous mesocarnivore in San Diego Bay. In 2019, they ranked third in Ecological Index – a product of having the second highest biomass despite only having the ninth highest numerical abundance. Like the Topsmelt and Round Stingray that rank higher, this species was ubiquitous throughout the bay during



these surveys, and were found in all sampling periods, ecoregions, depth strata and subhabitats. This important recreational fish species primarily utilizes bays and estuaries along the southern California coastline. Unlike previous years, there was no bimodal size distribution (Figure 17) during either sampling period and very few juveniles were present. Similar to Topsmelt, the April distribution is more typical of a July distribution, in this case representing either an extremely early reproductive season or a failed recruitment year for Spotted Sand Bass.




Figure 17. Total number of Spotted Sand Bass individuals by standard length (cm) from San Diego Bay, April and July 2019.

Slough Anchovy (*Anchoa delicatissima*)



Slough Anchovy ranked as the fourth most ecologically important fish species in San Diego Bay. This species was ubiquitous throughout the bay except for the North Ecoregion where it was not

captured during either survey. They were found in all depth strata and subhabitats. They only had the fifteenth highest biomass of all fishes captured in 2019 but were the second most abundant species captured during the surveys. Nearly all (99.8%) captured individuals were of adult size, with no change in size distribution between the April and July sampling periods (Figure 18). The bay is a well-known nursery area for this critical forage species and there were no doubt a significant number of juveniles of this species utilizing the bay, however juvenile Slough Anchovy are exceptionally narrow-bodied and are unlikely to be captured through most sampling methods employed.




Figure 18. Total number of Slough Anchovy individuals by standard length (cm) from San Diego Bay, April and July 2019.

Deepbody Anchovy (*Anchoa compressa*)

Deepbody Anchovy ranked fifth in Ecological Index. Despite being ranked so highly, this species was almost entirely limited to the Channel subhabitat during this study. In fact, 98.6% of the individuals



caught were from one purse seine replicate in the South-Central Ecoregion in July, and every individual was of adult size (Figure 19). Such a catch undoubtedly skews stock estimates and calculations of ecological importance, however as one of the rare species in the bay that almost exclusively inhabits the water column in the channel, it is difficult to assess their numbers accurately.




Figure 19. Total number of Deepbody Anchovy individuals by standard length (cm) from San Diego Bay, April and July 2019.

Giant Kelpfish (*Heterostichus rostratus*)



Giant Kelpfish ranked sixth in Ecological Index and were present during each sampling period in all ecoregions. They were found in both vegetated and non-vegetated habitats in the nearshore and intertidal depth strata, however, 15

times the number of individuals were taken in the nearshore strata (566) than the intertidal (39). 94.1% of the Giant Kelpfish captured were juveniles, and all individuals were less than 22 cm SL. There was a stark shift in distribution of size classes (Figure 20) from April to July suggesting a strong winter recruitment event, high juvenile mortality, and rapid growth between sampling events. This observed growth is consistent with that of previous surveys and fits within parameters for juvenile growth rates as estimated by a recent otolith ageing study (Winston et al. 2018).




Figure 20. Total number of Giant Kelpfish individuals by standard length (cm) from San Diego Bay, April and July 2019.

Shiner Perch (*Cymatogaster aggregata*)

Shiner Perch ranked seventh by the Ecological Index and were found during both sampling periods and at all ecoregions except the South Ecoregion in July. They were caught almost exclusively in the nearshore subhabitat. These mostly



juvenile fish (77.2%) were the sixth most abundant fish in the survey comprising 3.82% of the catch, but less than 1% of the biomass due to their small size. Their size frequency distributions (Figure 21) suggested a recruitment event prior to the April sampling efforts, and heavy natural mortality with slight growth between April and July. This distribution pattern is similar to that of other survey years.




Figure 21. Total number of Shiner Perch individuals by standard length (cm) from San Diego Bay, April and July 2019.

Kelp Pipefish (*Syngnathus californiensis*)



Though these long, narrow fishes only accounted for 0.16% of the biomass captured in 2019, Kelp Pipefish ranked eighth in

Ecological Index by accounting for 3.07% of the total abundance and being caught in all ecoregions during both survey periods. They were caught in all depth strata and habitats including a single individual in the channel subhabitat. More than 73% of the individuals were captured in the nearshore subhabitat and showed no apparent preference for vegetated versus non-vegetated habitat. A uniform size distribution belied the fact that nearly half (47.3%) of the individuals captured were juveniles (Figure 22), but the sharp increase in mean size from April to July supported the idea that this species grows quickly.




Figure 22. Total number of Kelp Pipefish individuals by standard length (cm) from San Diego Bay, April and July 2019.

California Halibut (*Paralichthys californicus*)

While not a frequently caught species (0.46% of the total catch), California Halibut ranked ninth in Ecological Index as a product of being caught at all ecoregions, all depth strata, and all subhabitats, and ranking eighth in biomass (1.7%). About 56% of the individuals were juveniles and only one fish would have been considered legal for recreational fishing purposes (Figure 23), but that fish alone weighed 2.5 kg – more than half of the total biomass of this species.




Figure 23. Total number of California Halibut individuals by standard length (cm) from San Diego Bay, April and July 2019.

Dwarf Perch (*Micrometrus minimus*)



Dwarf Perch ranked tenth in terms of Ecological Index, because it was the fourth most frequent catch (4.93%) with the tenth highest biomass (3.0%) during the 2019 surveys. It was only caught in the North Ecoregion, but during both survey periods, and in all subhabitats and depth strata except for the channel. As opposed to Shiner

Perch, this viviparous species appears to have reproduced sometime between the April and July sampling periods as there were far fewer adults and a far greater number of juveniles in July (Figure 24).




Figure 24. Total number of Dwarf Perch individuals by standard length (cm) from San Diego Bay, April and July 2019.

4.9 Catch by Sampling Method

The greatest number of species were collected in the purse seines (30 species), followed by large seine (23), beam trawl (21), otter trawl (15), small seine (11), and square enclosure (3; Tables 15 and 16). The purse seine captured the greatest number of fish, catching a total of 5,966, a number greatly influenced by large schools of anchovies and Topsmelt. There was an unusually large catch in the small seine (4,695), most of which were newly settled Topsmelt, and there were moderate catches in the large seine (1,610) and beam trawl (1,487), a relatively low catch in the otter trawl (256), and only 10 fish in the square enclosure (Table 15). The greatest amount of biomass was also captured in the purse seine (195.0 kg), with high biomass also captured in the otter trawl (36.4 kg) and beam trawl (29.6 kg). The large seine (9.14 kg) and small seine (1.30 kg) captured lower amounts of biomass, and the square enclosure captured only 20.5 g of fishes (Table 16).

The purse seine was most effective at sampling the schooling fishes (Slough Anchovy, Northern Anchovy, Topsmelt). The beam trawl was most effective for catching benthic nearshore and eelgrass fishes (Kelp Pipefish, Giant Kelpfish, Shiner Perch, Dwarf Perch). The square enclosure was largely ineffective in 2019 but was slightly more effective than in previous years. The beach seines were most effective at catching juvenile Topsmelt and Dwarf Perch. The top species caught in the otter trawls was Round Stingray. The highest density of fishes was captured in the small seine (1.578 individuals/m²) followed by the purse seine (0.280 individuals/m²; Table 17). The purse seines and beam trawls produced the highest biomass density values (9.151 g/m² and 2.126 g/m², respectively). The square enclosures captured the smallest amount of biomass (0.427 g/m²).



Transporting the square enclosure along the beach in the North-Central Ecoregion, July 2019.

Table 15. Total catch (number of individuals) of fish species taken in San Diego Bay in 2019 by sampling method.

PURSE SEINE				BEAM TRAWL			
Scientific Name	Common Name	Abundance	%	Scientific Name	Common Name	Abundance	%
<i>Atherinops affinis</i>	Topsmelt	2,563	42.96	<i>Heterostichus rostratus</i>	Giant Kelpfish	547	36.79
<i>Anchoa delicatissima</i>	Slough Anchovy	1,060	17.77	<i>Syngnathus californiensis</i>	Kelp Pipefish	303	20.38
<i>Anchoa compressa</i>	Deepbody Anchovy	1,032	17.30	<i>Micrometrus minimus</i>	Dwarf Perch	254	17.08
<i>Cymatogaster aggregata</i>	Shiner Perch	417	6.99	<i>Cymatogaster aggregata</i>	Shiner Perch	104	6.99
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	317	5.31	<i>Urobatis halleri</i>	Round Stingray	104	6.99
<i>Urobatis halleri</i>	Round Stingray	231	3.87	<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	52	3.50
<i>Micrometrus minimus</i>	Dwarf Perch	103	1.73	<i>Hypsoblennius gentilis</i>	Bay Blenny	31	2.08
<i>Sardinops sagax</i>	Pacific Sardine	78	1.31	<i>Paralichthys californicus</i>	California Halibut	22	1.48
<i>Leuresthes tenuis</i>	California Grunion	29	0.49	<i>Paralabrax nebulifer</i>	Barred Sand Bass	21	1.41
<i>Cynoscion parvipinnis</i>	Shortfin Corvina	20	0.34	<i>Clevelandia ios</i>	Arrow Goby	14	0.94
<i>Heterostichus rostratus</i>	Giant Kelpfish	19	0.32	<i>Porichthys myriaster</i>	Specklefin Midshipman	9	0.61
<i>Scomber japonicus</i>	Pacific Chub Mackerel	15	0.25	<i>Symphurus atricaudus</i>	California Tonguefish	9	0.61
<i>Haemulon californiensis</i>	Salema	14	0.23	<i>Halichoeres semicinctus</i>	Rock Wrasse	6	0.40
<i>Halichoeres semicinctus</i>	Rock Wrasse	13	0.22	<i>Paralabrax clathratus</i>	Kelp Bass	4	0.27
<i>Syngnathus californiensis</i>	Kelp Pipefish	12	0.20	<i>Acanthogobius flavimanus</i>	Yellowfin Goby	1	0.07
<i>Albula gilberti</i>	Cortez Bonefish	9	0.15	<i>Atherinops affinis</i>	Topsmelt	1	0.07
<i>Myliobatis californica</i>	Bat Ray	7	0.12	<i>Cheilotrema saturnum</i>	Black Croaker	1	0.07
<i>Paralabrax clathratus</i>	Kelp Bass	6	0.10	<i>Embiotoca jacksoni</i>	Black Perch	1	0.07
<i>Paralabrax nebulifer</i>	Barred Sand Bass	5	0.08	<i>Gymnura marmorata</i>	California Butterfly Ray	1	0.07
<i>Hippocampus ingens</i>	Pacific Seahorse	3	0.05	<i>Pleuronichthys decurrens</i>	Curlfin Sole	1	0.07
<i>Paralichthys californicus</i>	California Halibut	3	0.05	<i>Pleuronichthys ritteri</i>	Spotted Turbot	1	0.07
<i>Embiotoca jacksoni</i>	Black Perch	2	0.03				
<i>Dasyatis dipterura</i>	Diamond Stingray	1	0.02	# of Species:	21	1,487	
<i>Gibbonsia elegans</i>	Spotted Kelpfish	1	0.02				
<i>Hypsoblennius gentilis</i>	Bay Blenny	1	0.02				
<i>Phanerodon furcatus</i>	White Seaperch	1	0.02				
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	1	0.02				
<i>Porichthys myriaster</i>	Specklefin Midshipman	1	0.02				
<i>Seriphus politus</i>	Queenfish	1	0.02				
<i>Symphurus atricaudus</i>	California Tonguefish	1	0.02				
# of Species:	30	5,966					

Table 15 (continued).

LARGE SEINE		Abundance	%
Scientific Name	Common Name		
<i>Atherinops affinis</i>	Topsmelt	936	58.14
<i>Micrometrus minimus</i>	Dwarf Perch	297	18.45
<i>Anchoa delicatissima</i>	Slough Anchovy	205	12.73
<i>Heterostichus rostratus</i>	Giant Kelpfish	34	2.11
<i>Syngnathus californiensis</i>	Kelp Pipefish	23	1.43
<i>Clevelandia ios</i>	Arrow Goby	18	1.12
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	18	1.12
<i>Paralichthys californicus</i>	California Halibut	13	0.81
<i>Urobatis halleri</i>	Round Stingray	12	0.75
<i>Cymatogaster aggregata</i>	Shiner Perch	10	0.62
<i>Paralabrax nebulifer</i>	Barred Sand Bass	9	0.56
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin	8	0.50
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	5	0.31
<i>Embiotoca jacksoni</i>	Black Perch	4	0.25
<i>Hyporhamphus rosae</i>	California Halfbeak	4	0.25
<i>Hypsoblennius gentilis</i>	Bay Blenny	4	0.25
<i>Leuresthes tenuis</i>	California Grunion	3	0.19
<i>Fundulus parvipinnis</i>	California Killifish	2	0.12
<i>Anchoa compressa</i>	Deepbody Anchovy	1	0.06
<i>Atherinopsis californiensis</i>	Jacksmelt	1	0.06
<i>Ilypnus gilberti</i>	Cheekspot Goby	1	0.06
<i>Myliobatis californica</i>	Bat Ray	1	0.06
<i>Umbrina roncador</i>	Yellowfin Croaker	1	0.06
# of Species:	23	1,610	

SMALL SEINE		Abundance	%
Scientific Name	Common Name		
<i>Atherinops affinis</i>	Topsmelt	4,540	96.70
<i>Syngnathus californiensis</i>	Kelp Pipefish	92	1.96
<i>Micrometrus minimus</i>	Dwarf Perch	37	0.79
<i>Clevelandia ios</i>	Arrow Goby	15	0.32
<i>Heterostichus rostratus</i>	Giant Kelpfish	5	0.11
<i>Anchoa delicatissima</i>	Slough Anchovy	1	0.02
<i>Cymatogaster aggregata</i>	Shiner Perch	1	0.02
<i>Embiotoca jacksoni</i>	Black Perch	1	0.02
<i>Gymnura marmorata</i>	California Butterfly Ray	1	0.02
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin	1	0.02
<i>Paralichthys californicus</i>	California Halibut	1	0.02
# of Species:	11	4,695	

Table 15 (continued).

OTTER TRAWL			
Scientific Name	Common Name	Abundance	%
<i>Urobatis halleri</i>	Round Stingray	149	58.20
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	46	17.97
<i>Paralichthys californicus</i>	California Halibut	25	9.77
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	10	3.91
<i>Pleuronichthys ritteri</i>	Spotted Turbot	7	2.73
<i>Cymatogaster aggregata</i>	Shiner Perch	4	1.56
<i>Porichthys myriaster</i>	Specklefin Midshipman	3	1.17
<i>Cheilotrema saturnum</i>	Black Croaker	2	0.78
<i>Scorpaena guttata</i>	California Scorpionfish	2	0.78
<i>Symphurus atricaudus</i>	California Tonguefish	2	0.78
<i>Synodus lucioceps</i>	California Lizardfish	2	0.78
<i>Gymnura marmorata</i>	California Butterfly Ray	1	0.39
<i>Hippocampus ingens</i>	Pacific Seahorse	1	0.39
<i>Syngnathus californiensis</i>	Kelp Pipefish	1	0.39
<i>Xystreurus liolepis</i>	Fantail Sole	1	0.39
# of Species: 15		256	

SQUARE ENCLOSURE			
Scientific Name	Common Name	Abundance	%
<i>Clevelandia ios</i>	Arrow Goby	5	50.0
<i>Atherinops affinis</i>	Topsmelt	4	40.0
<i>Hypsoblennius gentilis</i>	Bay Blenny	1	10.0
# of Species: 3		10	

Table 16. Total biomass (g) of fish species taken from San Diego Bay in 2019 by sampling method.

PURSE SEINE				BEAM TRAWL			
Scientific Name	Common Name	Biomass (g)	%	Scientific Name	Common Name	Biomass (g)	%
<i>Urobatis halleri</i>	Round Stingray	68,380	35.06	<i>Urobatis halleri</i>	Round Stingray	18,195	61.47
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	53,959	27.67	<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	6,028	20.36
<i>Atherinops affinis</i>	Topsmelt	20,098	10.31	<i>Gymnura marmorata</i>	California Butterfly Ray	1,600	5.41
<i>Anchoa compressa</i>	Deepbody Anchovy	15,350	7.87	<i>Micrometrus minimus</i>	Dwarf Perch	1,459	4.93
<i>Myliobatis californica</i>	Bat Ray	8,400	4.31	<i>Heterostichus rostratus</i>	Giant Kelpfish	909	3.07
<i>Albula gilberti</i>	Cortez Bonefish	5,385	2.76	<i>Paralabrax nebulifer</i>	Barred Sand Bass	259	0.87
<i>Cynoscion parvipinnis</i>	Shortfin Corvina	5,108	2.62	<i>Cymatogaster aggregata</i>	Shiner Perch	221	0.74
<i>Dasyatis diptera</i>	Diamond Stingray	4,200	2.15	<i>Syngnathus californiensis</i>	Kelp Pipefish	213	0.72
<i>Paralichthys californicus</i>	California Halibut	2,527	1.30	<i>Halichoeres semicinctus</i>	Rock Wrasse	184	0.62
<i>Sardinops sagax</i>	Pacific Sardine	2,330	1.19	<i>Paralichthys californicus</i>	California Halibut	147	0.50
<i>Halichoeres semicinctus</i>	Rock Wrasse	1,654	0.85	<i>Porichthys myriaster</i>	Specklefin Midshipman	139	0.47
<i>Cymatogaster aggregata</i>	Shiner Perch	1,567	0.80	<i>Symphurus atricaudus</i>	California Tonguefish	80	0.27
<i>Anchoa delicatissima</i>	Slough Anchovy	1,440	0.74	<i>Hypsoblennius gentilis</i>	Bay Blenny	62	0.21
<i>Scomber japonicus</i>	Pacific Chub Mackerel	1,318	0.68	<i>Paralabrax clathratus</i>	Kelp Bass	35	0.12
<i>Paralabrax clathratus</i>	Kelp Bass	774	0.40	<i>Embiotoca jacksoni</i>	Black Perch	21	0.07
<i>Micrometrus minimus</i>	Dwarf Perch	639	0.33	<i>Pleuronichthys decurrens</i>	Curlfin Sole	21	0.07
<i>Leuresthes tenuis</i>	California Grunion	358	0.18	<i>Pleuronichthys ritteri</i>	Spotted Turbot	19	0.06
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	335	0.17	<i>Clevelandia ios</i>	Arrow Goby	5.5	0.02
<i>Porichthys myriaster</i>	Specklefin Midshipman	255	0.13	<i>Acanthogobius flavimanus</i>	Yellowfin Goby	3.0	0.01
<i>Haemulon californiensis</i>	Salema	190	0.10	<i>Atherinops affinis</i>	Topsmelt	1.5	0.01
<i>Phanerodon furcatus</i>	White Seaperch	190	0.10	<i>Cheilotrema saturnum</i>	Black Croaker	0.1	0.00
<i>Paralabrax nebulifer</i>	Barred Sand Bass	126	0.06				
<i>Heterostichus rostratus</i>	Giant Kelpfish	122	0.06	# of Species:	21	29,600	
<i>Syngnathus californiensis</i>	Kelp Pipefish	118	0.06				
<i>Hippocampus ingens</i>	Pacific Seahorse	105	0.05				
<i>Seriphus politus</i>	Queenfish	46	0.02				
<i>Embiotoca jacksoni</i>	Black Perch	23	0.01				
<i>Hypsoblennius gentilis</i>	Bay Blenny	11	0.01				
<i>Gibbonsia elegans</i>	Spotted Kelpfish	10	0.01				
<i>Symphurus atricaudus</i>	California Tonguefish	3	< 0.01				
# of Species:	30	195,021					

Table 16 (continued).

LARGE SEINE		Biomass	
Scientific Name	Common Name	(g)	%
<i>Urobatis halleri</i>	Round Stingray	2,729	29.86
<i>Atherinops affinis</i>	Topsmelt	1,887	20.64
<i>Micrometrus minimus</i>	Dwarf Perch	1,855	20.29
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	475	5.20
<i>Embiotoca jacksoni</i>	Black Perch	342	3.74
<i>Heterostichus rostratus</i>	Giant Kelpfish	334	3.65
<i>Anchoa delicatissima</i>	Slough Anchovy	290	3.17
<i>Myliobatis californica</i>	Bat Ray	265	2.90
<i>Paralabrax nebulifer</i>	Barred Sand Bass	223	2.44
<i>Paralichthys californicus</i>	California Halibut	208	2.28
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	100	1.09
<i>Atherinopsis californiensis</i>	Jacksmelt	66	0.72
<i>Cymatogaster aggregata</i>	Shiner Perch	65	0.71
<i>Leuresthes tenuis</i>	California Grunion	64	0.70
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin	57	0.62
<i>Hypsoblennius gentilis</i>	Bay Blenny	56	0.61
<i>Umbrina roncadore</i>	Yellowfin Croaker	49	0.54
<i>Syngnathus californiensis</i>	Kelp Pipefish	36	0.39
<i>Hyporhamphus rosae</i>	California Halfbeak	15	0.16
<i>Anchoa compressa</i>	Deepbody Anchovy	10	0.11
<i>Clevelandia ios</i>	Arrow Goby	8	0.09
<i>Fundulus parvipinnis</i>	California Killifish	4	0.04
<i>Ilypnus gilberti</i>	Cheekspot Goby	3	0.03
# of Species: 23		9,141	

SMALL SEINE		Biomass	
Scientific Name	Common Name	(g)	%
<i>Atherinops affinis</i>	Topsmelt	703	54.20
<i>Gymnura marmorata</i>	California Butterfly Ray	205	15.81
<i>Micrometrus minimus</i>	Dwarf Perch	181	13.96
<i>Embiotoca jacksoni</i>	Black Perch	115	8.87
<i>Syngnathus californiensis</i>	Kelp Pipefish	53	4.09
<i>Heterostichus rostratus</i>	Giant Kelpfish	23	1.77
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin	10	0.77
<i>Clevelandia ios</i>	Arrow Goby	4	0.27
<i>Anchoa delicatissima</i>	Slough Anchovy	2	0.15
<i>Cymatogaster aggregata</i>	Shiner Perch	1	0.08
<i>Paralichthys californicus</i>	California Halibut	1	0.04
# of Species: 11		1,297	

Table 16 (continued).

OTTER TRAWL			
Scientific Name	Common Name	Biomass	
		(g)	%
<i>Urobatis halleri</i>	Round Stingray	26,160	71.81
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	5,211	14.30
<i>Paralichthys californicus</i>	California Halibut	1,814	4.98
<i>Porichthys myriaster</i>	Specklefin Midshipman	982	2.70
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	849	2.33
<i>Pleuronichthys ritteri</i>	Spotted Turbot	495	1.36
<i>Gymnura marmorata</i>	California Butterfly Ray	300	0.82
<i>Synodus lucioceps</i>	California Lizardfish	255	0.70
<i>Scorpaena guttata</i>	California Scorpionfish	140	0.38
<i>Cheilotrema saturnum</i>	Black Croaker	135	0.37
<i>Symphurus atricaudus</i>	California Tonguefish	40	0.11
<i>Hippocampus ingens</i>	Pacific Seahorse	29	0.08
<i>Xystreureys liolepis</i>	Fantail Sole	10	0.03
<i>Cymatogaster aggregata</i>	Shiner Perch	6	0.02
<i>Syngnathus californiensis</i>	Kelp Pipefish	2	0.01
# of Species: 15		36,428	

SQUARE ENCLOSURE			
Scientific Name	Common Name	Biomass	
		(g)	%
<i>Hypsoblennius gentilis</i>	Bay Blenny	18	87.8
<i>Clevelandia ios</i>	Arrow Goby	1.5	7.3
<i>Atherinops affinis</i>	Topsmelt	1.0	4.9
# of Species: 3		20.5	

Table 17. Comparison of mean densities and biomass densities by gear type and sampling year for San Diego Bay. Values were calculated for years that were sampled in April and July and only includes data from those sampling months.

Year	Beam Trawl	Otter Trawl	Purse Seine	Large Seine	Small Seine	Square Enclosure	Year	Beam Trawl	Otter Trawl	Purse Seine	Large Seine	Small Seine	Square Enclosure		
Density (#/m ²)	1995	0.188	0.021	3.937	0.589	4.037	3.646	Biomass Density (g/m ²)	1995	3.455	2.229	10.44	0.816	0.527	1.681
	1996	0.170	0.009	8.205	0.658	8.699	6.396		1996	2.433	1.772	16.92	0.887	1.172	0.433
	1997	0.106	0.013	0.844	0.532	2.369	4.208		1997	1.170	1.844	6.195	1.504	0.300	0.409
	1998	0.037	0.003	1.995	0.427	0.403	4.000		1998	1.610	0.591	7.083	1.013	0.059	0.604
	2005	0.145	0.032	0.569	0.676	0.439	0.708		2005	5.137	1.426	5.580	1.684	0.217	0.196
	2008	0.223	0.006	0.390	0.171	0.702	0.542		2008	3.572	0.624	3.910	1.314	0.256	12.32
	2012	0.386	0.008	0.122	0.366	1.659	0.708		2012	7.199	1.026	7.949	1.502	1.044	2.065
	2015	0.184	0.028	0.705	0.219	0.869	0.146		2015	2.578	1.583	11.01	1.367	0.462	0.010
	2016	0.189	0.009	0.495	0.177	1.873	0.146		2016	2.188	1.288	6.356	0.764	0.390	0.019
	2019	0.107	0.004	0.280	0.152	1.578	0.208		2019	2.126	0.628	9.151	0.866	0.436	0.427

4.10 Best Estimates of Density and Standing Stock

The best estimate for the total stock size was 18,321,764 fishes (Table 18). With an estimated surface area of 4,858 ha (Table 2) this gives an overall fish density 0.38 individuals/m² (Table 18). The highest estimates were of Topsmelt (8.16 million), Deepbody Anchovy (2.75 million), Slough Anchovy (2.19 million), followed by Giant Kelpfish (1.09 million), and Shiner Perch (0.815 million). As is typical, schooling and forage fishes dominated the stock estimate for the bay.

The total best estimate of biomass standing stock was about 420 metric tons (MT) (Table 19). This gives an overall estimate of 8.64 g/m², about 35% higher than the 2016 estimate and above average for all historical surveys (Williams et al. 2016). The highest biomass estimates were of Round Stingray (136.5 MT), followed by Spotted Sand Bass (106.8 MT), Topsmelt (44.1 MT), Deepbody Anchovy (40.9 MT), and Bat Ray (17.0 MT). Round Stingray and Spotted Sand Bass are typically in the top five biomass estimates from past surveys, but Deepbody Anchovy is a historically atypical top biomass species.

Table 18. Best estimate of densities (#/m²) and stock estimates, April and July 2019.

Scientific Name	Common Name	Best Estimate of Density (#/m ²)				Weighted Mean	Stock Estimate (#)
		Depth Strata					
		Channel	Intertidal	Nearshore			
<i>Atherinops affinis</i>	Topsmelt	0.03913	1.52554	0.16082	0.16795	8,159,182	
<i>Anchoa compressa</i>	Deepbody Anchovy	0.14527	0.00009		0.05666	2,752,504	
<i>Anchoa delicatissima</i>	Slough Anchovy	0.01745	0.01941	0.06588	0.04513	2,192,640	
<i>Heterostichus rostratus</i>	Giant Kelpfish		0.00322	0.03930	0.02253	1,094,386	
<i>Cymatogaster aggregata</i>	Shiner Perch	0.00007	0.00095	0.02935	0.01679	815,856	
<i>Syngnathus californiensis</i>	Kelp Pipefish	0.00002	0.03091	0.02177	0.01365	663,147	
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	0.00183	0.00047	0.02140	0.01293	628,070	
<i>Micrometrus minimus</i>	Dwarf Perch		0.02813	0.01825	0.01153	559,926	
<i>Urobatis halleri</i>	Round Stingray	0.00257	0.00114	0.01555	0.00991	481,590	
<i>Clevelandia ios</i>	Arrow Goby		0.10417	0.00101	0.00474	230,266	
<i>Sardinops sagax</i>	Pacific Sardine	0.01098			0.00428	208,024	
<i>Hypsoblennius gentilis</i>	Bay Blenny	0.00014	0.02083	0.00223	0.00216	104,818	
<i>Leuresthes tenuis</i>	California Grunion		0.00028	0.00204	0.00117	57,071	
<i>Paralichthys californicus</i>	California Halibut	0.00043	0.00123	0.00158	0.00112	54,321	
<i>Cynoscion parvipinnis</i>	Shortfin Corvina	0.00239		0.00021	0.00105	51,185	
<i>Paralabrax nebulifer</i>	Barred Sand Bass		0.00085	0.00151	0.00089	43,431	
<i>Scomber japonicus</i>	Pacific Chub Mackerel	0.00211			0.00082	40,005	
<i>Haemulon californiensis</i>	Salema			0.00099	0.00056	27,285	
<i>Halichoeres semicinctus</i>	Rock Wrasse			0.00091	0.00052	25,336	
<i>Albula gilberti</i>	Cortez Bonefish	0.00099		0.00014	0.00046	22,567	
<i>Porichthys myriaster</i>	Specklefin Midshipman	0.00005		0.00065	0.00039	18,883	
<i>Symphurus atricaudus</i>	California Tonguefish	0.00003		0.00065	0.00038	18,557	
<i>Myliobatis californica</i>	Bat Ray	0.00028	0.00009	0.00035	0.00031	15,263	
<i>Paralabrax clathratus</i>	Kelp Bass			0.00042	0.00024	11,694	
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	0.00017	0.00170	0.00007	0.00018	8,527	
<i>Hippocampus ingens</i>	Pacific Seahorse	0.00002		0.00021	0.00013	6,173	
<i>Embiotoca jacksoni</i>	Black Perch		0.00038	0.00014	0.00010	4,634	
<i>Pleuronichthys ritteri</i>	Spotted Turbot	0.00012		0.00007	0.00009	4,276	
<i>Gymnura marmorata</i>	California Butterfly Ray	0.00002	0.00034	0.00007	0.00006	2,969	
<i>Cheilotrema saturnum</i>	Black Croaker	0.00003		0.00007	0.00005	2,642	
<i>Acanthogobius flavimanus</i>	Yellowfin Goby			0.00007	0.00004	1,989	
<i>Pleuronichthys decurrens</i>	Curlfin Sole			0.00007	0.00004	1,989	
<i>Dasyatis diptera</i>	Diamond Stingray			0.00007	0.00004	1,949	
<i>Gibbonsia elegans</i>	Spotted Kelpfish			0.00007	0.00004	1,949	
<i>Phanerodon furcatus</i>	White Seaperch			0.00007	0.00004	1,949	
<i>Seriphus politus</i>	Queenfish			0.00007	0.00004	1,949	
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin		0.00076		0.00003	1,472	
<i>Hyporhamphus rosae</i>	California Halfbeak		0.00038		0.00002	736	
<i>Scorpaena guttata</i>	California Scorpionfish	0.00003			0.00001	653	
<i>Synodus lucioceps</i>	California Lizardfish	0.00003			0.00001	653	
<i>Fundulus parvipinnis</i>	California Killifish		0.00019		0.00001	368	
<i>Xystreurus liolepis</i>	Fantail Sole	0.00002			0.00001	327	
<i>Atherinopsis californiensis</i>	Jacksmelt		0.00009		< 0.00001	184	
<i>Ilypnus gilberti</i>	Cheekspot Goby		0.00009		< 0.00001	184	
<i>Umbrina roncadior</i>	Yellowfin Croaker		0.00009		< 0.00001	184	
Grand Totals:		0.22420	1.74135	0.38606	0.37713	18,321,764	

Table 19. Best estimate of biomass densities (g/m²) and standing stock, April and July 2019.

Scientific Name	Common Name	Best Estimate of Density (g/m ²)				Weighted Mean	Stock Estimate (kg)	Stock Estimate (MT)
		Depth Strata						
		Channel	Intertidal	Nearshore				
<i>Urobatis halleri</i>	Round Stingray	0.45097	0.25843	4.60318	2.81003	136,511	136.5	
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	0.30039	0.04498	3.64759	2.19808	106,783	106.8	
<i>Atherinops affinis</i>	Topsmelt	0.87725	0.23622	0.97593	0.90786	44,104	44.1	
<i>Anchoa compressa</i>	Deepbody Anchovy	2.16075	0.00095		0.84273	40,940	40.9	
<i>Myliobatis californica</i>	Bat Ray	0.10557	0.02509	0.53843	0.34908	16,958	17.0	
<i>Albula gilberti</i>	Cortez Bonefish	0.56799		0.09502	0.27568	13,392	13.4	
<i>Cynoscion parvipinnis</i>	Shortfin Corvina	0.38007		0.16948	0.24483	11,894	11.89	
<i>Dasyatis dipterura</i>	Diamond Stingray			0.29561	0.16850	8,186	8.19	
<i>Sardinops sagax</i>	Pacific Sardine	0.32798			0.12791	6,214	6.21	
<i>Paralichthys californicus</i>	California Halibut	0.03127	0.01970	0.17786	0.11436	5,556	5.56	
<i>Scomber japonicus</i>	Pacific Chub Mackerel	0.18553			0.07236	3,515	3.52	
<i>Gymnura marmorata</i>	California Butterfly Ray	0.00517	0.06888	0.11494	0.07029	3,415	3.41	
<i>Micrometrus minimus</i>	Dwarf Perch		0.17566	0.10481	0.06677	3,244	3.24	
<i>Halichoeres semicinctus</i>	Rock Wrasse			0.11641	0.06636	3,224	3.22	
<i>Cymatogaster aggregata</i>	Shiner Perch	0.00010	0.00616	0.11029	0.06315	3,068	3.07	
<i>Anchoa delicatissima</i>	Slough Anchovy	0.02590	0.02746	0.08840	0.06159	2,992	2.99	
<i>Heterostichus rostratus</i>	Giant Kelpfish		0.03163	0.06527	0.03847	1,869	1.87	
<i>Paralabrax clathratus</i>	Kelp Bass			0.05448	0.03105	1,508	1.51	
<i>Pleuronichthys guttulatus</i>	Diamond Turbot	0.01464	0.00947	0.02358	0.01953	949	0.95	
<i>Hypsoblennius gentilis</i>	Bay Blenny	0.00155	0.37500	0.00445	0.01814	881	0.88	
<i>Porichthys myriaster</i>	Specklefin Midshipman	0.01693		0.01795	0.01683	818	0.82	
<i>Leuresthes tenuis</i>	California Grunion		0.00606	0.02520	0.01460	709	0.71	
<i>Paralabrax nebulifer</i>	Barred Sand Bass		0.02112	0.01861	0.01145	556	0.56	
<i>Syngnathus californiensis</i>	Kelp Pipefish	0.00003	0.01781	0.01527	0.00943	458	0.46	
<i>Haemulon californiensis</i>	Salema			0.01337	0.00762	370	0.37	
<i>Phanerodon furcatus</i>	White Seaperch			0.01337	0.00762	370	0.37	
<i>Hippocampus ingens</i>	Pacific Seahorse	0.00050		0.00739	0.00441	214	0.21	
<i>Pleuronichthys ritteri</i>	Spotted Turbot	0.00853		0.00136	0.00411	199	0.20	
<i>Symphurus atricaudus</i>	California Tonguefish	0.00069		0.00575	0.00354	172	0.17	
<i>Embiotoca jacksoni</i>	Black Perch		0.03864	0.00162	0.00247	120	0.12	
<i>Seriphus politus</i>	Queenfish			0.00324	0.00185	90	0.09	
<i>Synodus lucioceps</i>	California Lizardfish	0.00440			0.00171	83	0.08	
<i>Clevelandia ios</i>	Arrow Goby		0.03125	0.00040	0.00148	72	0.07	
<i>Scorpaena guttata</i>	California Scorpionfish	0.00241			0.00094	46	0.05	
<i>Cheilotrema saturnum</i>	Black Croaker	0.00233		0.00001	0.00091	44	0.04	
<i>Pleuronichthys decurrens</i>	Curlfin Sole			0.00151	0.00086	42	0.04	
<i>Gibbonsia elegans</i>	Spotted Kelpfish			0.00070	0.00040	19	0.02	
<i>Atherinopsis californiensis</i>	Jacksmelt		0.00625		0.00025	12	0.01	
<i>Leptocottus armatus</i>	Pacific Staghorn Sculpin		0.00540		0.00022	10	0.01	
<i>Umbrina roncadore</i>	Yellowfin Croaker		0.00464		0.00019	9	0.01	
<i>Acanthogobius flavimanus</i>	Yellowfin Goby			0.00022	0.00012	6	0.01	
<i>Xystreurus liolepis</i>	Fantail Sole	0.00017			0.00007	3	< 0.01	
<i>Hyporhamphus rosae</i>	California Halfbeak		0.00142		0.00006	3	< 0.01	
<i>Fundulus parvipinnis</i>	California Killifish		0.00038		0.00002	1	< 0.01	
<i>Ilypnus gilberti</i>	Cheekspot Goby		0.00028		0.00001	1	< 0.01	
Grand Totals:		5.47115	1.41288	11.31169	8.63793	419,630	419.6	

4.11 Avian Forage Species

Forage species are primarily surface-dwelling schooling fish that are accessible to diving avian predators, especially terns. Generally, forage fishes are small silvery-sided fishes that are found in large schools. These schooling fishes are generally not habitat specific and move throughout the bay's ecosystem. Thirteen species of important forage fishes (as defined in Pondella and Williams 2011) were captured during this study. The most abundant forage fishes were Topsmelt, Slough Anchovy, and Deepbody Anchovy. Topsmelt were primarily found at small (juvenile) size classes (<50 mm SL) appropriate for nesting birds to feed their young in the area. The typical timing for the recruitment of fishes to San Diego Bay begins in the spring and continues through the summer, but it appears this began much earlier in the year in 2019. The biomass standing stock estimate for forage fish was 106.7 MT. When estimating by ecoregion, values were highest at the South-Central Ecoregion (66.5 MT) which was driven almost entirely by one catch of Deepbody Anchovy. This was followed by the North Ecoregion (32.3 MT), the North-Central Ecoregion (14.0 MT), and the South Ecoregion (3.4 MT; Table 20).



Surf scoter (*Melanitta perspicillata*) swimming near Sweetwater Marsh.

Table 20. Best estimate of biomass standing stock for forage fish species by ecoregion, 2019.

Ecoregion	Depth Strata					Weighted Mean	Stock Estimate (kg)	Stock Estimate (MT)
	Scientific Name	Common Name	Channel	Intertidal	Nearshore			
North	<i>Atherinops affinis</i>	Topsmelt	2.87162	0.49530	2.33587	2.55224	25,063	25.1
	<i>Scomber japonicus</i>	Pacific Chub Mackerel	0.67568			0.41216	4,047	4.05
	<i>Micrometrus minimus</i>	Dwarf Perch		0.70265	0.41925	0.18051	1,773	1.77
	<i>Cymatogaster aggregata</i>	Shiner Perch		0.01061	0.19088	0.06363	625	0.62
	<i>Heterostichus rostratus</i>	Giant Kelpfish		0.09697	0.16868	0.06148	604	0.60
	<i>Leuresthes tenuis</i>	California Grunion		0.02424	0.04505	0.01632	160	0.16
	<i>Clevelandia ios</i>	Arrow Goby			0.00057	0.00019	2	< 0.01
	Grand Total:		1.32977	3.16030	3.28654	32,274	32.3	
North-Central	<i>Atherinops affinis</i>	Topsmelt	0.59122	0.43616	1.42652	0.90088	7,279	7.28
	<i>Sardinops sagax</i>	Pacific Sardine	1.23874			0.70608	5,705	5.71
	<i>Cymatogaster aggregata</i>	Shiner Perch		0.01061	0.11064	0.04257	344	0.34
	<i>Scomber japonicus</i>	Pacific Chub Mackerel	0.06644			0.03787	306	0.31
	<i>Heterostichus rostratus</i>	Giant Kelpfish		0.02727	0.05402	0.02189	177	0.18
	<i>Leuresthes tenuis</i>	California Grunion			0.05574	0.02118	171	0.17
	<i>Anchoa delicatissima</i>	Slough Anchovy		0.05492	0.00169	0.00339	27	0.03
	<i>Clevelandia ios</i>	Arrow Goby			0.00086	0.00033	3	< 0.01
	<i>Anchoa compressa</i>	Deepbody Anchovy		0.00379		0.00019	2	< 0.01
	Grand Total:	1.89640	0.53275	1.64948	1.73439	14,014	14.0	
South-Central	<i>Anchoa compressa</i>	Deepbody Anchovy	8.55856			3.08108	61,776	61.78
	<i>Anchoa delicatissima</i>	Slough Anchovy		0.05492	0.15709	0.09748	1,954	1.95
	<i>Cymatogaster aggregata</i>	Shiner Perch		0.00152	0.07658	0.04676	937	0.94
	<i>Atherinops affinis</i>	Topsmelt		0.04167	0.07123	0.04470	896	0.90
	<i>Sardinops sagax</i>	Pacific Sardine	0.07320			0.02635	528	0.53
	<i>Heterostichus rostratus</i>	Giant Kelpfish		0.00076	0.03089	0.01887	378	0.38
	<i>Clevelandia ios</i>	Arrow Goby		0.08333	0.00014	0.00259	52	0.05
	<i>Atherinopsis californiensis</i>	Jacksmelt		0.02500		0.00075	15	0.02
	Grand Total:	8.63176	0.20720	0.33593	3.31857	66,537	66.5	
South	<i>Anchoa delicatissima</i>	Slough Anchovy	0.10360		0.19482	0.17517	1,864	1.86
	<i>Atherinops affinis</i>	Topsmelt	0.04617	0.14242	0.07010	0.06988	744	0.74
	<i>Cymatogaster aggregata</i>	Shiner Perch	0.00041	0.00189	0.06306	0.05247	558	0.56
	<i>Anchoa compressa</i>	Deepbody Anchovy	0.08446			0.01098	117	0.12
	<i>Heterostichus rostratus</i>	Giant Kelpfish		0.00152	0.00760	0.00637	68	0.07
	<i>Clevelandia ios</i>	Arrow Goby		0.04167		0.00167	18	0.02
	<i>Hyporhamphus rosae</i>	California Halfbeak		0.00568		0.00023	2	0.00
	<i>Fundulus parvipinnis</i>	California Killifish		0.00152		0.00006	1	0.00
	Grand Total:	0.23465	0.19470	0.33559	0.31683	3,371	3.4	

4.12 Fisheries Species

During this study, 15 species were captured that have importance in either the recreational or commercial fisheries in California. The most abundant fisheries species were Spotted Sand Bass, Pacific Sardine, and California Halibut. Including all ecoregions, standing stock estimates of fisheries species totaled 149.7 MT. Like the forage fishes, when estimating by ecoregion values were greatest at the South-Central Ecoregion (71.4 MT), this time driven by Spotted Sand Bass, Cortez Bonefish, and Shortfin Corvina. Stock estimates were next highest in the South Ecoregion (93.2 MT) and the North-Central Ecoregion (26.6 MT), both driven almost exclusively by Spotted Sand Bass. The North Ecoregion had the smallest stock estimate at 13.0 MT (Table 21).

Table 21. Best estimate of biomass standing stock for recreational/commercial fishery species by ecoregion, 2019.

Ecoregion	Scientific Name	Common Name	Depth Strata			Weighted Mean	Stock Estimate (kg)	Stock Estimate (MT)
			Channel	Intertidal	Nearshore			
North	<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	0.04244		1.54279	0.53501	5,254	5.3
	<i>Scomber japonicus</i>	Pacific Chub Mackerel	0.67568			0.41216	4,047	4.05
	<i>Paralichthys californicus</i>	California Halibut	0.08254		0.70383	0.28261	2,775	2.78
	<i>Paralabrax clathratus</i>	Kelp Bass			0.21791	0.07191	706	0.71
	<i>Embiotoca jacksoni</i>	Black Perch		0.15457	0.00648	0.01141	112	0.11
	<i>Scorpaena guttata</i>	California Scorpionfish	0.00965			0.00589	58	0.06
	<i>Paralabrax nebulifer</i>	Barred Sand Bass		0.00720		0.00043	4	< 0.01
	Grand Total:		0.81031	0.16177	2.47100	1.31943	12,957	13.0
North-Central	<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	0.01931	0.03333	6.33305	2.41923	19,547	19.55
	<i>Sardinops sagax</i>	Pacific Sardine	1.23874			0.70608	5,705	5.71
	<i>Cynoscion parvipinnis</i>	Shortfin Corvina			0.19707	0.07489	605	0.61
	<i>Scomber japonicus</i>	Pacific Chub Mackerel	0.06644			0.03787	306	0.31
	<i>Paralichthys californicus</i>	California Halibut	0.01358	0.01250	0.03851	0.02300	186	0.19
	<i>Paralabrax nebulifer</i>	Barred Sand Bass		0.06742	0.05086	0.02270	183	0.18
	<i>Cheilotrema saturnum</i>	Black Croaker	0.00931			0.00531	43	0.04
	<i>Umbrina roncador</i>	Yellowfin Croaker		0.01856		0.00093	7	0.01
	<i>Paralabrax clathratus</i>	Kelp Bass			0.00115	0.00044	4	< 0.01
	Grand Total:	1.34738	0.13182	6.62064	3.29044	26,587	26.6	
South-Central	<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	0.62387	0.10227	2.47128	1.73515	34,790	34.79
	<i>Albula gilberti</i>	Cortez Bonefish	1.99606		0.38007	0.95042	19,056	19.06
	<i>Cynoscion parvipinnis</i>	Shortfin Corvina	1.52027		0.45045	0.82207	16,483	16.48
	<i>Sardinops sagax</i>	Pacific Sardine	0.07320			0.02635	528	0.53
	<i>Paralabrax nebulifer</i>	Barred Sand Bass		0.00985	0.02356	0.01467	294	0.29
	<i>Paralichthys californicus</i>	California Halibut	0.02565	0.03561	0.00169	0.01133	227	0.23
	<i>Cheilotrema saturnum</i>	Black Croaker			0.00003	0.00002	0	< 0.01
	Grand Total:	4.23905	0.14773	3.32708	3.56001	71,378	71.4	
South	<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	0.57770	0.04432	4.24324	3.59877	38,291	38.29
	<i>Albula gilberti</i>	Cortez Bonefish	0.27590			0.03587	382	0.38
	<i>Cynoscion parvipinnis</i>	Shortfin Corvina			0.03041	0.02524	269	0.27
	<i>Paralabrax nebulifer</i>	Barred Sand Bass			0.01436	0.01192	127	0.13
	<i>Seriphus politus</i>	Queenfish			0.01295	0.01075	114	0.11
	<i>Paralichthys californicus</i>	California Halibut	0.00331	0.03068		0.00166	18	0.02
	Grand Total:	0.85691	0.07500	4.30096	3.68419	39,200	39.2	

4.13 Southern (Panamic) Species Found in San Diego Bay

San Diego Bay is known for being the northern edge of the range for many southern fishes that are not normally distributed in the Southern California Bight. As an example, at least 25 northern range extensions have been reported from the bay (Table 22). During this study, six species [Cortez Bonefish, California Butterfly Ray (*Gymnura marmorata*), Pacific Seahorse (*Hippocampus ingens*), California Halfbeak (*Hyporhamphus rosae*), Diamond Stingray (*Dasyatis dipterura*), and the largest Shortfin Corvina (48 cm SL) captured in these surveys] with primarily southern distributions were taken (Table 23). These fishes were found almost exclusively in the southern half of the bay and none were captured in the North Ecoregion.

Table 22. Panamic species previously recorded in San Diego Bay.

Scientific Name	Common Name	First Recorded SDB Collection Date	Citation
<i>Albula gilberti</i>	Cortez Bonefish	prior to 1918	Starks (1918)
<i>Caranx caballus</i>	Green Jack	1857	Girard (1858)
<i>Caranx caninus</i>	Pacific Crevalle Jack	16 Mar 1972	Miller and Lea (1972)
<i>Caranx vinctus</i>	Cocinero	12 Aug 1997	Lea and Rosenblatt (2000)
<i>Caranx sexfasciatus</i>	Bigeye Trevally	Nov 1990	Lea and Walker (1995)
<i>Cetengraulis mysticetus</i>	Anchoveta	1980-1986	Duffy (1987)
<i>Chanos chanos</i>	Milkfish	22 Mar 1982	Duffy and Bernard (1985)
<i>Chaetodon humeralis</i>	Threebanded Butterflyfish	1857	Girard (1858)
<i>Ctenogobius sagittula</i>	Longtail Goby	1907	Lea and Rosenblatt (2000)
<i>Cynoscion parvipinnis</i>	Shortfin Corvina	common	Jordan and Gilbert (1880)
<i>Dasyatis dipterura</i>	Diamond Stingray	1880 (type locale)	Jordan and Gilbert (1880)
<i>Elops affinis</i>	Machete	30 Dec 1997	Lea and Rosenblatt (2000)
<i>Gymnura marmorata</i>	California Butterfly Ray	1864 (type locale)	Cooper (1864)
<i>Haemulon flaviguttatum</i>	Cortez Grunt	May 1991	Lea and Rosenblatt (1992)
<i>Hippocampus ingens</i>	Pacific Seahorse	1855 (type locale)	Girard (1858)
<i>Hyporhamphus rosae</i>	California Halfbeak	1880 (type locale)	Jordan and Gilbert (1880)
<i>Lobotes pacificus</i>	Pacific Tripletail	5 Nov 1997	Lea and Rosenblatt (2000)
<i>Mugil curema</i>	White Mullet	25 May 1985	Lea et al. (1988)
<i>Polydactylus approximans</i>	Blue Bobo	Dec 1997	Lea and Rosenblatt (2000)
<i>Pseudupeneus grandisquamous</i>	Bigscale Goatfish	15 Jul 1998	Lea and Rosenblatt (2000)
<i>Scomberomorus sierra</i>	Pacific Sierra	Dec 1995	Williams et al. (2011)
<i>Selene brevoorii</i>	Mexican Lookdown	Feb 1993	Lea and Walker (1995)
<i>Sphyrna lewini</i>	Scalloped Hammerhead	Apr 1997	Lea and Rosenblatt (2000)
<i>Strongylura exilis</i>	California Needlefish	common	Fitch and Lavenberg (1975)
<i>Zapteryx exasperata</i>	Banded Guitarfish	1880 (type locale)	Jordan and Gilbert (1880)



Pacific Seahorses caught by purse seine in the South Ecoregion in July 2019.

Table 23. Abundance of Panamic species collected in San Diego Bay by ecoregion, April and July 2019.

Scientific Name	Common Name	Ecoregions							
		North		North-Central		South-Central		South	
		April	July	April	July	April	July	April	July
<i>Albula gilberti</i>	Cortez Bonefish					5	3	1	
<i>Cynoscion parvipinnis</i>	Shortfin Corvina			1			18		1
<i>Dasyatis dipterura</i>	Diamond Stingray							1	
<i>Gymnura marmorata</i>	California Butterfly Ray							3	
<i>Hippocampus ingens</i>	Pacific Seahorse							4	
<i>Hyporhamphus rosae</i>	California Halfbeak							4	

4.14 Indigenous Bay and Estuary Fishes

As the largest estuary in southern California, San Diego Bay provides critical habitat for bay and estuary fishes and continues to function as a nursery area for nearly two-thirds (65.2%) of those fishes. The high productivity rate coupled with the abundance of juvenile fishes in the bay highlights the importance of the bay as a nursery habitat. The bay contains extensive shallow water eelgrass habitat that supports a unique assemblage of juvenile and adult fishes that, in turn, support surrounding nearshore ecosystems. Juvenile fishes emigrate from the bay to offshore habitats, and important or endangered avian species utilize forage fishes in the bay. Southern California indigenous bay and estuary fishes represented 23.1% of the total catch in this survey (Table 24).

Table 24. Indigenous bay/estuarine species taken in San Diego Bay by ecoregion in 2019.

Scientific Name	Common Name	Ecoregions				Total	%
		North	North-Central	South-Central	South		
<i>Anchoa delicatissima</i>	Slough Anchovy	110	475	681		1,266	39.05
<i>Anchoa compressa</i>	Deepbody Anchovy	1	1,019	13		1,033	31.86
<i>Syngnathus californiensis</i>	Kelp Pipefish	65	121	163	82	431	13.29
<i>Paralabrax maculatofasciatus</i>	Spotted Sand Bass	155	81	147	37	420	12.95
<i>Clevelandia ios</i>	Arrow Goby	5	18	21	8	52	1.60
<i>Hypsoblennius gentilis</i>	Bay Blenny	17			20	37	1.14
<i>Fundulus parvipinnis</i>	California Killifish			2		2	0.06
<i>Ilypnus gilberti</i>	Cheekspot Goby			1		1	0.03
Total % of catch:		5.7%	42.8%	46.0%	9.5%		23.1%

4.15 Invasive Species

In addition to being a warm-water refuge for southern species, San Diego Bay is also a major port-of-entry and commercial shipping hub. Releases of ballast water and trans-Pacific transportation of hull fouling organisms and their associated hitchhikers into historically disturbed habitat provides ideal opportunities for invasive species, such as Yellowfin Goby (*Acanthogobius flavimanus*) and Chameleon Goby (*Tridentiger trigonocephalus*) to establish themselves in the bay. Though ballast water exchange in the bay has been regulated to help reduced the threat since 2000, these species were established prior to the implementation of these regulations. The Yellowfin Goby was first described inside tidal marshes of the South Ecoregion by Williams et al. (1998) and has been reported in many brackish and freshwater areas in California where they pose a threat to native fish species as predators. Although the low-salinity requirements of this species appear to limit its expansion potential, no eradication or control efforts for this invasive have been successful (Molnar et al. 2008). Williams et al. (1998) recommended management actions that reduce off-season freshwater inflows and return tidal action to impounded saltmarsh areas in order to favor native species and prevent further spread of exotics.

The Chameleon Goby was first captured in San Diego Bay in January 1995 during the Allen et al. (2002) survey, and subsequently described with additional records by Pondella and Chinn (2005). Despite the possibility of competing with native species for habitat, this invader has not become enough of a problem to require management action, and there are no known natural controls in California's marine environment (Molnar et al. 2008). Ironically, the Chameleon Goby may be controlled by Yellowfin Goby predation (Meng et al. 1994).

During the 2012 survey, sampling yielded both of those species: three Yellowfin Gobies and 18 Chameleon Gobies. Given the widespread nature of Chameleon Goby throughout the bay during those surveys (captured in the channel, nearshore vegetated, and nearshore non-vegetated areas, in all ecoregions but the North), we reported that there may be a sustained invasion and self-recruiting population of Chameleon Goby within the bay (Williams and Pondella 2012). However, no Chameleon Gobies were caught in 2015, 2016, or 2019 (Williams et al. 2015, Williams et al. 2016). The only Yellowfin Goby encountered in 2015 was a single partially digested individual that was regurgitated by a Spotted Sand Bass in the South Ecoregion. Two Yellowfin Gobies were caught in the South Ecoregion in 2016, and only one was captured in 2019 in the North Ecoregion. The paucity of Yellowfin and Chameleon Gobies may be a product of heavy predation, low reproductive success, or simply more effective filtering of ballast water. Rainfall in Winter 2018-2019 likely produced favorable conditions for both of these species as both species thrive in fresh to brackish water environments. However, any advantage was not noticeable given the infrequent capture of either species.

4.16 Comparison of the Current and Historical April and July Surveys

4.16.1 Abundance, Biomass, and Stock Estimates

Total catch and biomass from the April and July 2019 sampling periods were also compared to values from 1995-1998, 2005, 2008, 2012, 2015, and 2016. While the 2019 surveys represent the fewest fishes ever captured in historical surveys, the biomass captured was only slightly below average (Tables 25-28; Figures 25-28). This is a product of not capturing large numbers of small individuals as is typical during April sampling, but instead capturing fewer, but larger, fish throughout the survey. This was particularly true in the South Ecoregion where fewer than 30% of the historical mean number of individuals was captured, but the biomass captured was the second highest ever recorded – more than 30% higher than average. Consequently, the 2019 stock estimate was less than 20% of the mean number of individuals while the biomass standing stock estimate was nearly average (98.8% of the mean; Table 29).

Table 25. Total abundance by sampling year. Results were calculated for years that were sampled in April and July and only includes data from those sampling months.

Ecoregion	Sampling Years									
	1995	1996	1997	1998	2005	2008	2012	2015	2016	2019
North	59,178	91,176	8,978	14,486	4,237	7,233	4,244	10,240	5,158	6,238
North-Central	19,523	112,964	8,718	11,603	12,539	3,354	5,645	5,874	7,932	4,005
South-Central	22,403	3,623	10,659	8,267	2,346	2,666	3,422	4,789	3,001	2,233
South	5,063	3,153	4,735	14,738	5,337	2,438	3,952	3,240	5,036	1,548
Total:	106,167	210,916	33,090	49,094	24,459	15,691	17,263	24,143	21,127	14,024




Figure 25. Total abundance by ecoregion and sampling year. Results were calculated for years that were sampled in April and July and only includes data from those sampling months.

Table 26. Total abundance by sampling year. Results were calculated for years that were sampled in April and July and only includes data from those sampling months.

Common Name	Sampling Years									
	1995	1996	1997	1998	2005	2008	2012	2015	2016	2019
Northern Anchovy	52,389	147,173	3	8,373	1,397	10	0	6,666	1,205	0
Topsmelt	25,272	33,915	12,917	9,232	7,448	3,393	6,529	2,024	4,136	8,044
Slough Anchovy	16,821	16,616	11,029	27,790	11,219	5,538	1,566	7,957	8,825	1,266
Shiner Perch	3,998	3,222	2,794	336	1,180	1,700	2,174	915	199	536
California Grunion	0	739	3,136	1,123	186	0	0	1,608	965	32
Arrow Goby	401	339	104	73	62	927	2,438	629	1,749	52
Deepbody Anchovy	69	221	29	5	2	72	17	80	30	1,033
Other	7,217	8,691	3,078	2,162	2,965	4,051	4,539	4,264	4,018	3,061




Figure 26. Total abundance of the most frequently caught species by sampling year. Results were calculated for years that were sampled in April and July and only includes data from those sampling months.

Table 27. Total biomass (kg) of fishes captured during April and July surveys by ecoregion.

Ecoregion	Sampling Years									
	1995	1996	1997	1998	2005	2008	2012	2015	2016	2019
North	111.9	195.4	70.1	58.7	58.9	36.5	119.7	112.8	83.2	69.0
North-Central	97.2	192.3	88.4	74.4	121.0	55.3	83.0	120.9	65.3	55.1
South-Central	111.8	46.6	65.4	33.2	34.2	43.7	70.7	69.2	49.2	59.7
South	89.2	75.8	48.2	52.3	77.4	49.0	74.8	75.3	52.2	87.7
Total:	410.0	510.1	272.1	218.6	291.6	184.5	348.2	378.2	249.9	271.5




Figure 27. Total biomass (kg) of fishes captured during April and July surveys by ecoregion.

Table 28. Total biomass (kg) of the highest biomass fishes captured during April and July surveys by species and sampling year.

Common Name	Sampling Years									
	1995	1996	1997	1998	2005	2008	2012	2015	2016	2016
Round Stingray	79.1	63.5	77.2	25.9	109.8	60.9	134.2	129.9	110.9	115.5
Spotted Sand Bass	59.9	29.7	33.5	42.2	36.3	47.3	62.3	50.8	28.2	65.7
Northern Anchovy	49.7	178.9	0.0	9.5	3.0	0.0	0.0	50.9	29.0	0.0
Slough Anchovy	45.7	30.1	19.3	66.0	17.4	10.2	2.4	14.7	14.8	1.7
Topsmelt	22.9	20.5	44.4	20.5	12.1	10.3	11.4	4.7	4.6	22.7
Bat Ray	26.3	37.2	0.0	0.5	36.5	0.0	31.3	26.2	4.6	8.7
Barred Sand Bass	38.3	40.2	10.6	4.7	4.7	2.7	2.2	5.6	7.5	0.6
California Butterfly Ray	0.0	0.0	0.0	8.6	2.1	5.1	15.5	51.2	2.4	2.1
Other	88.2	109.9	87.1	40.7	69.7	48.0	88.9	44.1	47.9	54.6




Figure 28. Total biomass (kg) of the highest biomass fishes captured during April and July surveys, shown by species over time (left) and as a proportion of the annual catch (right).

Table 29. Stock estimates and biomass standing stock by sampling year. Estimates were calculated for years that were sampled in April and July and only includes data from those sampling months.

Survey Year (source)	Stock Estimate (#)	Standing Stock (kg)
1995 (Allen et al. 2002)	178,106,064	513,340
1996 (Allen et al. 2002)	368,102,566	769,855
1997 (Allen et al. 2002)	49,326,442	302,962
1998 (Allen et al. 2002)	101,099,343	349,989
2005 (Pondella et al. 2006)	31,258,743	343,308
2008 (Pondella and Williams 2009a)	24,805,106	249,398
2012 (Williams and Pondella 2012)	16,180,679	465,376
2015 (Williams et al. 2015)	35,545,986	518,885
2016 (Williams et al. 2016)	30,173,603	311,227
2019 (Present Study)	18,321,764	419,630

4.16.2 Community Metrics

Shannon Diversity (H') and species richness values were determined for April and July from the previous surveys (Allen 1999, Pondella et al. 2006, Pondella and Williams 2009a, Williams and Pondella 2012, Williams et al. 2015, Williams et al. 2016) to allow direct comparisons of the data sets. The 1995-1998 survey years were used for the comparison because these were the only years from Allen et al. (2002) where both April and July were sampled. Overall, 2019 H' estimates were variable among ecoregions and rank among historical values. The North and North-Central Ecoregions had the third and second lowest diversity values, respectively, of any sampling year. However, diversity values for the South-Central and South Ecoregions were the third highest of any sampling year (Table 26, Figure 25). Species richness for 2019 was the lowest ever recorded in both the North and North-Central Ecoregions and was tied for third lowest at both the South-Central and South Ecoregions for any survey period (Table 27; Figure 26).



R/V Old Blue heading to the South-Central Ecoregion in April 2019.

Table 30. Shannon diversity (H') values by sampling year. Estimates were calculated for years that were sampled in April and July and only includes data from those sampling months.

Ecoregion	Sampling Years									
	1995	1996	1997	1998	2005	2008	2012	2015	2016	2019
North	0.74	0.90	1.34	1.42	1.77	1.72	1.56	1.43	1.94	0.93
North-Central	1.75	0.93	1.50	1.25	1.36	1.62	1.63	1.81	1.61	1.18
South-Central	1.32	1.72	1.13	0.37	1.77	1.88	1.92	1.41	1.51	1.77
South	1.93	1.84	1.35	0.59	1.06	2.03	1.84	1.71	1.61	1.82
Total:	1.46	1.04	1.65	1.31	1.65	2.05	2.02	2.05	2.00	1.69




Figure 29. Shannon diversity (H') values by ecoregion and sampling year. Estimates were calculated for years that were sampled in April and July and only includes data from those sampling months.

Table 31. Species richness values by sampling year. Estimates were calculated for years that were sampled in April and July and only includes data from those sampling months.

Ecoregion	Sampling Years									
	1995	1996	1997	1998	2005	2008	2012	2015	2016	2019
North	35	42	29	34	38	33	30	35	40	28
North-Central	32	33	31	26	38	27	37	35	33	23
South-Central	31	23	27	22	25	23	32	27	21	22
South	34	24	26	30	23	25	29	23	31	24
Total:	53	54	42	51	57	48	52	52	55	45




Figure 30. Species richness values by ecoregion and sampling year. Estimates were calculated for years that were sampled in April and July and only includes data from those sampling months.

4.16.3 Community Structure

Six significantly different groups (a-f) were determined by cluster analysis of fish communities for each survey period (Figure 31). The driving species behind each community group can be visualized in Figure 32, where circles are scaled to mean abundance by community group of eleven typical San Diego Bay fish taxa. Sampling periods prior to, during, and after the 1997-1998 El Niño event showed the strongest differentiation from most other surveys. The April and July 1997 sampling periods form their own distinct fish community (group ‘f’), driven by comparatively higher catch of Pacific Sardine and California Grunion. After substantial rainfall during the winter of 1997-1998, the catch in April 1998 (group ‘a’) was dominated by Topsmelt, but also had large catches of Striped Mullet (*Mugil cephalus*) and Cortez Bonefish, two estuarine residents that thrive in brackish water. Relatively little else was captured during that survey. By July 1998, the fish community returned to a typical mid-1990’s July pattern (group ‘d’) where Northern Anchovy (*Engraulis mordax*) dominated the catch.




Figure 31. Fish communities for the twenty-two April and July surveys were clustered in this tree. Branch distances shown in this diagram represent the relative similarity of stations to each other using Bray-Curtis distances and group-average linkage. Significantly different (SIMPROF, $\alpha = 0.05$) fish communities are indicated by letters (a-f) and color groupings.

In spite of the intensity of the 2015-2016 El Niño event, all four sampling periods (April 2015-July 2016) clustered into the largest group ('b'), as did both April and July of 2019. Community structure of fishes in 2019 was most similar to that of the 2012 surveys but was not statistically different than any other survey performed in the 21st Century.




Figure 32. Relative mean abundance (circles scaled to value) per sampling period for select fish taxa in San Diego Bay by community group (as identified in Figure 30).

5.0 Literature Cited

- Allen, L. G. 1980. Seasonal abundance, composition, and productivity of the littoral fish assemblage in upper Newport Bay, California. U.S. Fish Bull., 80(4): 769-790.
- Allen, L. G., A. M. Findlay, and C. M. Phalen. 2002. Structure and standing stock of the fish assemblages of San Diego Bay, California from 1994-1999. Bull. So. Calif. Acad. Sci. 101(2), 49-85.
- Allen, L. G. 1999. Fisheries inventory and utilization of San Diego Bay, San Diego, California. Final report for contract to the U.S. Navy Naval Engineering Naval Command Southwest Division and the San Diego Unified Port District, 138 pp.
- Clarke K. R., P. J. Somerfield, R. N. Gorley. 2008. Testing of null hypotheses in exploratory community analyses: similarity profiles and biota-environment linkage. J. Exp. Mar. Biol. Ecol. 366: 56-69.
- Cooper, J.G. 1864. On new genera and species of California fishes. No. III. Proc. Cal. Acad. Sci. Ser. 1, v. 3 (sigs. 7-8): 108-114.
- Duffy, J. M. 1987. A review of the San Diego Bay striped mullet, *Mugil cephalus*, fishery. Calif. Dept. Fish Game, Mar. Res. Tech. Rep. No. 56, 10 pp.
- Duffy, J. M. and H. J. Bernard. 1985. Milkfish, *Chanos chanos* (Forsskal, 1775), taken in southern California adds new family (Channidae) to the California marine fauna. Calif. Fish Game, 71(2): 122-125.
- Girard, C. F. 1858. Fishes. In: U.S. War Department, Reports of explorations and surveys, to ascertain the most practicable and economical route for a railroad from the Mississippi River to the Pacific Ocean, v. 10, part 4 (Washington, D.C.)
- Jordan, D. S. and C. H. Gilbert. 1880. Notes on a collection of fishes from San Diego, California. Proc. U. S. Nat. Mus., 3:23-34
- Jordan, D. S., B. W. Evermann and H. W. Clark. 1930. Check list of the fishes and fish-like vertebrates of North and Middle America north of the northern boundary of Venezuela and Columbia. Appendix X to the Report of the United States Commission of Fish and Fisheries for 1928. Washington, DC: Government Printing Office. 670 pp.
- Lea, R. N., C. C. Swift, and R. J. Lavenberg. 1988. Records of *Mugil curema* Valenciennes, the white mullet, from southern California. Bull. So. Calif. Acad. Sci., 87(1): 31-34.
- Lea, R. N. and R. H. Rosenblatt. 1992. The Cortez grunt (*Haemulon flaviguttatum*) recorded from two embayments in southern California. Calif. Fish Game, 78(4): 163-165.
- Lea, R. N. and H. J. Walker, Jr. 1995. Record of the bigeye trevally, *Caranx sexfasciatus*, and Mexican lookdown, *Selene brevoorti*, with notes on other carangids from California. Calif. Fish Game, 81(3): 89-95.
- Lea, R. N. and R. H. Rosenblatt. 2000. Observations on fishes associated with the El Niño off California. CalCOFI Rep. 41: 117-129.
- Meng, L., P. B. Moyle, and B. Herbold. 1994. Changes in abundance and distribution of native and introduced fishes of Suisun Marsh. Trans. Am. Fish. Soc., 123(4):498-507

- Molnar, J. L., R. L. Gamboa, C. Revenga, and M. D. Spalding. 2008. Assessing the global threat of invasive species to marine biodiversity. *Front. Ecol. Environ.* 6(9): 485-492
- Oksanen J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R. B. O'Hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, H. Wagner. 2019. *vegan: Community Ecology Package*. R package version 2.5-5. <http://CRAN.R-project.org/package=vegan>
- Peeling, T. J. 1975. A proximate biological survey of San Diego Bay, California. No. NUC-TP-389-REV-1. Naval Undersea Center, San Diego, CA, Report No. TP389.
- Pondella, D. J., II, J. Froeschke and B. Young. 2006. Fisheries Inventory and Utilization of San Diego Bay, San Diego California for surveys conducted in April and July 2005. February 2006. 103 p.
- Pondella, D. J., II and J. P. Williams. 2009a. Fisheries Inventory and Utilization of San Diego Bay, San Diego California for surveys conducted in April and July 2008. February 2009. 68 p.
- Pondella, D. J., II and J. P. Williams. 2009b. Fisheries Inventory and Utilization of San Diego Bay, San Diego California for surveys conducted in June 2009. June 2009. 24 p.
- Pondella, D. J., II and J. P. Williams. 2011. Summary and analysis of past fish collection data, with comparison to past California Least Tern productivity, San Diego: oceanographic indices, forage fish and tern breeding success. March 2011. 249 p.
- R Core Team. 2019. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>
- San Diego County MS4 Co-permittees. 2008. San Diego Bay Watershed Urban Runoff Management Program Document. Final, March 2008. Submitted to the San Diego Regional Water Quality Control Board by the San Diego Bay Copermittees.
- Shannon, C. E. 1948. A mathematical theory of communication. *Bell System Technical Journal*, 27, 379-423.
- Starks, E. C. 1918. The herrings and herring-like fishes of California. *Calif. Fish and Game* 4(2): 58-65.
- Wang, P. F., R.T, Cheng, K. Richter, E. S. Gross, D. Sutton, J.W. Gartner. 1998. Modeling tidal hydrodynamics of San Diego Bay, California. *Journal of the American Water Resources Association*, 34(5): 1123-1140.
- Whitaker D., Christman M. 2014. *clustsig: Significant Cluster Analysis*. R package version 1.1. <http://CRAN.R-project.org/package=clustsig>
- Williams, G. D., J. S. Desmond, and J. B. Zedler. 1998. Extension of two nonindigenous fishes, *Acanthogobius flavimanus* and *Poecilia latipinna*, into San Diego Bay marsh habitats. *Calif. Fish Game* 84(1): 1-17.
- Williams, J. P., D. J. Pondella II, B. M. Haggin, and L. G. Allen. 2011. New record of Pacific sierra (*Scomberomorus sierra*) with notes on previous California records. *Calif. Fish Game* 97(1): 43-46.
- Williams, J. P. and D. J. Pondella II. 2012. Fisheries Inventory and Utilization of San Diego Bay, San Diego California for surveys conducted in April and July 2012. September 2012. 66 p.

- Williams, J. P., D. J. Pondella II, C. M. Williams, S. Schwab. 2015. Fisheries Inventory and Utilization of San Diego Bay, San Diego California for surveys conducted in April and July 2015. September 2015. 64 p.
- Williams, J. P., D. J. Pondella II, C. M. Williams, M. J. Robart. 2016. Fisheries Inventory and Utilization Study to Determine Impacts from El Niño in San Diego Bay, San Diego California for surveys conducted in April and July 2016. November 2015. 68 p.
- Winston, M., J. T. Claisse, C. M. Williams, J. P. Williams, D. J. Pondella II. 2018. Age and growth of the Giant Kelpfish, *Heterostichus rostratus* Girard 1854, in southern California, USA. *Journal of Applied Ichthyology* 34(1): 91-96.



Sam Soule ('18) with a Cortez Bonefish caught by purse seine in the South Ecoregion in April 2019.